

# The bHLH Transcription Factor NeuroD Governs Photoreceptor Genesis and Regeneration Through Delta-Notch Signaling

Scott M. Taylor,<sup>1</sup> Karen Alvarez-Delfin,<sup>2</sup> Carole J. Saade,<sup>2</sup> Jennifer L. Thomas,<sup>3</sup> Ryan Thummel,<sup>3</sup> James M. Fadool,<sup>2</sup> and Peter F. Hitchcock<sup>1</sup>

<sup>1</sup>Department of Ophthalmology and Visual Sciences, University of Michigan, W. K. Kellogg Eye Center, Ann Arbor, Michigan, United States

<sup>2</sup>Department of Biological Science, Florida State University, Tallahassee, Florida, United States

<sup>3</sup>Departments of Anatomy/Cell Biology and Ophthalmology, Wayne State University, Detroit, Michigan, United States

Correspondence: Peter F. Hitchcock, University of Michigan, Department of Ophthalmology and Visual Sciences, W. K. Kellogg Eye Center, 1000 Wall Street, Ann Arbor, MI 48105, USA; peterh@med.umich.edu.

Submitted: July 3, 2015

Accepted: October 6, 2015

Citation: Taylor SM, Alvarez-Delfin K, Saade CJ, et al. The bHLH transcription factor NeuroD governs photoreceptor genesis and regeneration through Delta-Notch signaling. *Invest Ophthalmol Vis Sci.* 2015;56:7496-7515. DOI:10.1167/iovs.15-17616

**PURPOSE.** Photoreceptor genesis in the retina requires precise regulation of progenitor cell competence, cell cycle exit, and differentiation, although information around the mechanisms that govern these events currently is lacking. In zebrafish, the basic helix-loop-helix (bHLH) transcription factor NeuroD governs photoreceptor genesis, but the signaling pathways through which NeuroD functions are unknown. The purpose of this study was to identify these pathways, and during photoreceptor genesis, Notch signaling was investigated as the putative mediator of NeuroD function.

**METHODS.** In embryos, genetic mosaic analysis was used to determine if NeuroD functions is cell- or non-cell-autonomous. Morpholino-induced NeuroD knockdown, CRISPR/Cas9 mutation, and pharmacologic and transgenic approaches were used, followed by in situ hybridization, immunocytochemistry, and quantitative RT-PCR (qRT-PCR), to identify mechanisms through which NeuroD functions. In adults, following photoreceptor ablation and NeuroD knockdown, similar methods as above were used to identify NeuroD function during photoreceptor regeneration.

**RESULTS.** In embryos, NeuroD function is non-cell-autonomous, NeuroD knockdown increases Notch pathway gene expression, Notch inhibition rescues the NeuroD knockdown-induced deficiency in cell cycle exit but not photoreceptor maturation, and Notch activation and CRISPR/Cas9 mutation of *neurod* recapitulate NeuroD knockdown. In adults, NeuroD knockdown prevents cell cycle exit and photoreceptor regeneration and increases Notch pathway gene expression, and Notch inhibition rescues this phenotype.

**CONCLUSIONS.** These data demonstrate that during embryonic development, NeuroD governs photoreceptor genesis via non-cell-autonomous mechanisms and that, during photoreceptor development and regeneration, Notch signaling is a mechanistic link between NeuroD and cell cycle exit. In contrast, during embryonic development, NeuroD governs photoreceptor maturation via mechanisms that are independent of Notch signaling.

**Keywords:** photoreceptor development, photoreceptor regeneration, rods, cones, cell cycle

The vertebrate retina is composed of seven cell types, generated in a conserved sequence from a pool of multipotent, mitotic progenitors.<sup>1</sup> Several basic helix-loop-helix (bHLH) transcription factors have prominent roles in the retina to regulate cell fate specification,<sup>2-5</sup> cell cycle exit, and neuronal differentiation and maturation.<sup>6,7</sup> NeuroD is a proneural bHLH transcription factor identified in vivo as a neuronal determination factor,<sup>8</sup> and in vitro as a link between cell cycle exit and neuronal differentiation.<sup>9</sup> In the mouse retina, NeuroD is required for survival of rods<sup>10</sup> and regulates opsin selection among cones.<sup>11</sup> In the chick, NeuroD is required for photoreceptor development and survival.<sup>12</sup> In medaka fish, NeuroD and Six6 coregulate each other to control amacrine cell and photoreceptor genesis.<sup>6</sup> In embryonic zebrafish, *neurod* is expressed in mitotic photoreceptor progenitors,<sup>13,14</sup> and this expression is regulated by the zinc

finger protein, *Insm1a*.<sup>15</sup> Within the photoreceptor lineages NeuroD governs cell cycle exit and photoreceptor maturation.<sup>7</sup>

The Notch pathway mediates cell-to-cell communication through receptor-ligand interactions. Notch receptors are expressed on the cell surface and interact with membrane-bound ligands (e.g., Delta, Jagged), regulating transcription in apposing cells.<sup>16,17</sup> In vertebrate retinal development, Notch signaling regulates the balance between neurogenesis and gliogenesis,<sup>18-20</sup> maintains progenitors in an undifferentiated, proliferative state,<sup>19,21</sup> specifies cell fates, and governs the onset of neurogenesis.<sup>22</sup> These events can be regulated in the retina through transcriptional control of Notch signaling molecules. For example, in the chick and mouse, the bHLH transcription factor *Ascl1a* governs cell cycle exit and differentiation through regulation of the Notch ligand *dll*,<sup>23,24</sup> and in zebrafish the HLH-binding cofactor *Id2a* regulates the

transition from proliferation to differentiation through regulation of Notch ligands and receptors.<sup>21</sup>

Following retinal damage in zebrafish, Müller glia function as intrinsic stem cells that dedifferentiate, reenter the cell cycle, and produce neurogenic progenitors that regenerate lost neurons.<sup>25–33</sup> Recent studies in zebrafish have identified mechanisms that govern the Müller glia response to retinal injury,<sup>34–37</sup> but information is lacking around the pathways that govern photoreceptor regeneration from Müller glia-derived progenitor cells. The mechanisms that govern retinal regeneration are expected to largely recapitulate embryonic development,<sup>28,29,38</sup> though few molecules have been functionally studied during both events. NeuroD function has not been studied in the adult zebrafish retina, but following photoreceptor ablation *neurod* is expressed in Müller glia-derived mitotic progenitors,<sup>39</sup> suggesting that NeuroD has a role in photoreceptor regeneration.

The goal of the current study is to identify the mechanisms that govern photoreceptor genesis from the pool of multipotent progenitors in the embryo and from stem cell-derived progenitors in the adult, by elucidating the pathways through which NeuroD functions during photoreceptor development and regeneration, respectively. In embryos, reciprocal transplant chimeric analysis shows that for cell cycle exit and photoreceptor maturation, NeuroD function is non-cell-autonomous. Knockdown of NeuroD and CRISPR/Cas9 targeted mutation of *neurod* prevent cell cycle exit and photoreceptor maturation, and increase expression of Notch pathway molecules. Inhibition of Notch signaling rescues deficiencies in cell cycle exit but not photoreceptor maturation. In adults, NeuroD knockdown prevents cell cycle exit among injury-induced progenitors and photoreceptor regeneration, and this, too, is rescued by Notch inhibition. These data demonstrated a conserved function for NeuroD during photoreceptor genesis and regeneration, and identified Notch signaling as a molecular mechanism that links these events.

## METHODS

These studies adhered to the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research and were approved by the University of Michigan Institutional Animal Care and Use Committee.

### NeuroD KnockDown in Embryos

We used AB strain zebrafish (*Danio rerio*) for developmental experiments and they were purchased from the Zebrafish International Research Center (ZIRC; University of Oregon, Portland, OR, USA). Embryos were collected within 15 minutes of spawning and incubated at 28.5°C on a 14/10-hour light/dark cycle. Morpholino oligonucleotides (MO; Gene Tools, LLC, Philomath, OR, USA) were used to induce NeuroD knockdown. The *neurod* MO (ATG 5'-TGACTTCGTCATGTCG GAACTCTAG-3') and *neurod* MM control MO (5'-TGAGTTGGT CATCTCGCAACTGTAG-3') have been described previously.<sup>7</sup> Morpholino oligonucleotides were diluted in 1 × Danieau buffer<sup>40</sup> and 5 ng MOs were injected at the 1 cell stage.

### Systemic Labeling With 5-Bromo-2'-Deoxyuridine (BrdU) or 5-Ethynyl-2'-Deoxyuridine (EdU)

Cells in S-phase of the cell cycle were labeled with either BrdU or EdU. Embryos were incubated for 20 minutes in ice-cold (48 hours post fertilization [hpf]) or room-temperature (70 hpf) 10 mM BrdU or 1.5 mM EdU, dissolved in embryo rearing solution containing 15% dimethylsulfoxide (DMSO). For BrdU staining,

sections were incubated in 100°C sodium citrate buffer (10 mM sodium citrate, 0.05% Tween 20, pH 6.0) for 30 minutes to denature DNA and cooled at room temperature for 20 minutes. Sections then were subjected to standard immunolabeling as described below. Ethynyl deoxyuridine was visualized using the Click-it EdU kit (Invitrogen, Carlsbad, CA, USA).

## Chimeric Analysis

Chimeric mosaic analysis was performed at the shield stage as previously described.<sup>41–43</sup> Reciprocal transplants were performed between wild-type (WT) and NeuroD knockdown embryos and larvae were incubated in EdU as described above. To label donor cells, embryos were injected with 5% tetramethyl rhodamine-dextran (D-1816, 10,000 MW; Life Technologies, Carlsbad, CA, USA) at the 1- to 2-cell stage. To evaluate the differentiation of rod photoreceptors, donor cells were taken from Tg(XOPS:GFP) embryos, in which green fluorescent protein (GFP) is driven by promoter elements of *Xenopus* rhodopsin.<sup>44</sup> To evaluate cone differentiation, retinal sections were immunostained for markers of cone photoreceptors as described below.

Mosaic analysis data were collected from larvae at 72 to 75 hpf. Retinas containing a narrow radial clone of donor-derived cells were selected for analysis. Confocal images were collected using a Leica SP5 confocal microscope (Leica, Wetzlar, Germany), and data were taken from 5- $\mu$ m thick Z-stacks through clones of donor-derived cells. Within each clone, the number of dextran-labeled cells within the outer nuclear layer was counted and scored for the presence of EdU, GFP, or the cone cell marker, Zpr-1. Five clones of donor cells were evaluated per treatment group.

## CRISPR/Cas9 Targeted Mutation of *neurod*

The sgRNA target sequence for *neurod* (GGACGACGAGGAA GAAGAAGAGG) was identified using ZiFiT software (available in the public domain at [zifit.partners.org](http://zifit.partners.org)). Cas9 mRNA and sgRNA were generated according to previously described methods.<sup>45</sup> Single cell-stage embryos were injected with 1 nL solution containing 100 pg/nL sgRNA and 150 pg/nL Cas9 mRNA. Embryos were collected for histology and genetic sequencing at 70 hpf, following incubation in EdU as described above. Embryo heads were collected and fixed in 4% buffered paraformaldehyde for histology. For each individual embryo, the hindbrain and tails were digested at 55°C for 3 hours in PCR extraction buffer containing 10 mM Tris-HCl (pH8), 2 mM EDTA, 0.2% Triton X-100, and 200  $\mu$ g/mL Proteinase K.<sup>46</sup> Samples then were heated to 95°C, centrifuged, and stored at –20°C.

## T7 Assay for Detection of CRISPR-Induced Insertions or Deletions

The T7 endonuclease assay was conducted according to established protocols (available in the public domain at [www.crisprflydesign.org](http://www.crisprflydesign.org)). Briefly, screening primers (F: 5'-GCGCTCTAGTGCATTTTCATC-3', R: 5'-TAGTTCTTGGCT AGTCGGAG) were designed to amplify a 530 bp region around the *neurod* sgRNA target site. We performed PCR using standard protocols for Platinum Taq-HF polymerase (Invitrogen), using 10  $\mu$ L PCR extraction buffer (above) as the DNA template. Purified DNA product (200 ng) was used for the analysis, and 10 U T7 endonuclease I (New England Biolabs, Ipswich, MA, USA) was used for the digest. The presence of CRISPR-induced insertions or deletions (indels) was indicated by a double band approximately 250 to 300 bp. For embryos that were positive for indels, the PCR product was subcloned

into the pGEM-T Easy vector (Promega, Madison, WI, USA), and 6 to 9 clones were sequenced for each embryo. Mutations were detected using a pairwise blast (National Center for Biotechnology Information [NCBI], Bethesda, MD, USA) against WT DNA and amino acid sequences determined using the ExPASy translate tool (available in the public domain at [www.expasy.org](http://www.expasy.org)).

### Inhibition of Notch Signaling With DAPT

The Notch pathway was inhibited with the  $\gamma$ -secretase inhibitor DAPT (N-[N-(3,5-Difluorophenacetyl-L-alanyl)]-S-phenylglycine *t*-Butyl Ester, 565784; Calbiochem, EMD Millipore, Billerica, MA, USA), which inhibits Notch cleavage and activation of the intracellular domain.<sup>47</sup> DAPT was diluted in embryo (E3) media for a working concentration of 50  $\mu$ M (in 0.2% DMSO).<sup>21</sup> Embryos were treated from 48 to 70 hpf and exposed to systemic BrdU (described above) for 20 minutes before being killed. Control embryos were treated in E3 with a matched concentration of DMSO only.

To quantify BrdU- and/or EdU-labeled cells, for each 10- $\mu$ m-thick section, a confocal z-stack was collected through the entire section using a Leica SP5 confocal microscope. Each section contained optic nerve to standardize the retinal location analyzed in each embryo. Cells were quantified in the entire z-stack using the Spots function of Leica Imaris Software with standardized parameters for sensitivity and cell size. Following DAPT treatment in embryos, BrdU-positive cells were counted in one section from one eye of each embryo ( $n = 8$  embryos per treatment). Statistical significance was determined using a Student's *t*-test.

### Transgenic Lines and Conditional Activation of the Notch Pathway

The tol2 pT2KhsppGGFF vector (generous gift from Koichi Kawakami, National Institute of Genetics, Mishima, Shizuoka Japan) containing a GFP-Gal4-FF fusion protein (GGFF) downstream of the hsp70 promoter<sup>48</sup> was injected into WT zebrafish embryos at the 1-cell stage (~25–50 pg) along with tol2 transposon mRNA (~25–50 pg). F0 embryos were screened for GFP expression in the lens at 72 hpf, raised, and outcrossed to WT adults, and F1 embryos were screened in the same way. Hemizygous F1 Tg(hsp70:GGFF) adults then were crossed with hemizygous Tg(UAS:*myc-notch1a:intra*).<sup>49</sup> Embryos were heat-shocked in a 38°C water bath for 1 hour at 52 to 53 hpf, and screened for GFP expression at 69 hpf. GFP-positive embryos were selected for analysis, indicating that they carried the GGFF transgene. Of these, 50% carried Tg(hsp70:GGFF) alone and were used as controls, and 50% carried both transgenes, confirmed post hoc by immunolabeling with anti-c-Myc. Embryos were incubated in 1.5 mM EdU for 20 minutes (described above) immediately before being killed at 70 hpf. For quantification of EdU+ cells in the ONL, confocal z-stacks were collected through one entire section of a single eye per embryo ( $n = 8$  Myc+,  $n = 5$  Myc– [WT]). Only sections passing through the optic nerve were used as a means to standardize the retinal location analyzed. EdU+ cells within the ONL were quantified as described above.

### Histology, Immunohistochemistry, and In Situ Hybridization

For histology, tissues were fixed overnight at 4°C in phosphate buffered 4% paraformaldehyde, infiltrated with 20% sucrose, then frozen in optical cutting temperature (OCT) medium. Immunostaining and in situ hybridization were performed on 10- $\mu$ m-thick cross-sections through the central retina in the

vicinity of the optic nerve. Immunostaining was performed using previously published protocols.<sup>50</sup> The primary and secondary antibodies and dilutions used here were: mouse anti-BrdU 1:100 (347580; BD Biosciences, Franklin Lakes, NJ, USA), mouse anti-PCNA 1:100 (8825; Sigma-Aldrich Corp., St. Louis, MO, USA), Zpr-1 1:200 (anti-Arrestin 3, red-green double cones, ZIRC), Zpr-3 1:200 (anti-Rhodopsin, rods, ZIRC), anti-c-Myc 1:200 (9E10; ThermoFisher Scientific, Halethorp, MD, USA), rabbit anti-GFP (ab290; Abcam, Cambridge, UK), and goat anti-mouse Alexa Fluor 488 and 555 1:500 (Life Technologies). Following immunohistochemistry, when applicable, a TUNEL labeling In situ Cell Death Detection Kit (Roche Diagnostic Corp., Indianapolis, IN, USA) was used to compare cell death between treatment groups.

In situ hybridizations were performed using previously published protocols.<sup>13,51</sup> Riboprobes for *ascl1a*, *notch1a*, *ber4*, *deltaA*, and *deltaD* were generated from PCR products,<sup>52</sup> and riboprobes for *rhodopsin*, *pde6c*, *neurod*, *crx*, *nr2e3*, and *pax6a* were generated from full-length cDNA clones. In vitro transcription was performed using the appropriate polymerase and DIG or fluorescein labeling kit (Roche Diagnostic Corp.). Double in situ hybridization for *notch1a* and *dla* was performed as described previously.<sup>13</sup> For comparisons of relative expression, tissue sections from control and experimental groups were placed on the same slide and developed for identical periods of time.

### Reverse Transcriptase Quantitative Real-Time PCR (qRT-PCR)

Total RNA was extracted from embryo forebrains and eyes (heads) using the Aurum Total RNA Mini Kit and following the manufacturer's protocol (Bio-Rad Laboratories, Inc., Hercules, CA, USA). Three to five biological replicates were collected for each treatment and each time point (30–50 heads per replicate). Reverse transcription was performed using the Qiagen QuantiTect Reverse Transcription kit according to the manufacturer's instructions (Qiagen, Venlo, The Netherlands). Each biological replicate was run in triplicate using 6 ng cDNA and IQ SYBR Green Supermix (Bio-Rad Laboratories, Inc.) and a Bio-Rad 384 Well Real Time PCR machine. Primer sets are listed in Table 1. Expression was calculated from  $\Delta$ CT values and normalized to *bactin* expression using Bio-Rad CFX Manager Software. Fold changes were calculated by dividing normalized expression by control values. Statistical significance was determined using a Student's *t*-test.

### Photoreceptor Ablation in Adults

The Tg(*neurod:egfp*) zebrafish line<sup>53</sup> was used for regeneration experiments and was previously shown to be an accurate reporter for endogenous *neurod* expression.<sup>39</sup> Adult fish were housed in recirculating systems at 28.5°C on a 14/10-hour light/dark cycle. To ablate retinal photoreceptors, fish were exposed to bright light as described previously.<sup>54–57</sup> For the NeuroD knockdown experiments in which proliferating progenitors and regenerated photoreceptors were counted (see below), *albino* fish were used to consistently ablate all photoreceptors in the dorsal retina.<sup>57</sup> For experiments in which colocalization of EGFP and other molecules were analyzed using immunocytochemistry and in situ hybridization, complete ablation of photoreceptors was not required and, therefore, fully pigmented fish were used. The regeneration response is identical in *albino* and pigmented fish (Taylor SM, Hitchcock PF, unpublished data, 2014). Before bright light exposure, *albino* fish were housed in complete darkness for 10 days and pigmented fish for 24 hours. Immediately after dark adaptation, fish were exposed to high intensity light (ca.

TABLE 1. Primer Sets Used for qRT-PCR

Gene	Forward Primer 5'-	Reverse Primer 5'-
<i>cyclinD1</i>	CAAACACGCCAGACCTTTGTG	GGGTCACTTCTGATGACTTGCG
<i>cyclinB1</i>	AGTTTAGGCTGCTTCAGGAGAC	ATGCACGGTCTGTACAAAAGC
<i>ascl1a</i>	GGGCTCATAACGACCCCTCTGA	TCCAAGCGAGTGCTGATATTT
<i>notch1a</i>	GTCGTGCTGTTGAAGAATGGAGC	GGTCCAGGTGATCTGCAATATCC
<i>ber4</i>	CACTGGATCAATCAGCAGCA	TCTCGGCGGATCTTCTCC
<i>neuroD</i>	AGACACACATCCACGCGCATCTCC	TCCGTTGAGACCGGCATCATCGTC
<i>bes6</i>	CGAAGGAGGCTGACAGTGTG	ATTCGAGGAGGTGGTTCAGC
<i>deltaA</i>	GATCGACCACTGTTCTTCCA	TTGGCAAGGGTACATCGAAC
<i>deltaB</i>	CGGCGAGGCTTGTTCAGATTAC	CCTGGATAATGAACGCACTCGC
<i>deltaC</i>	ACGCCGCTGGCAACAGAATTTG	GCAAGTGCCTGAAAAGGCTGG
<i>deltaD</i>	GGTGCTCGGATCAAACCTCTTC	ATGCGACTGATCAGACGGTCTG
<i>six6a</i>	CTGGACTCACACCCACGCAAGTG	GACTGACTTCTGGGCTTGAGGCG
<i>bactin</i>	GCAATGAGCGTTTCCGTTG	TGTGTTGGCATAACAGTCC

100,000 lux) from a mercury arc lamp for 30 minutes and then to constant light (ca. 20,000 lux) for 72 hours.<sup>56,57</sup> Finally, to analyze expression of molecules in Müller glia, the Tg(*gfa-p:egfp*)<sup>58</sup> fish line and EGFP immunocytochemistry were used to identify Müller glia cell bodies and processes. For all experiments, days post lesion (dpl) indicates the number of days following the onset of light exposure.

### Adult Morpholino Electroporation

For adult experiments, *neurod* or standard control MO (5'-CTCTTACCTCAGTTACAATTTATA-3') were purchased from Gene Tools, LLC with a 3' lissamine (sulforhodamine B) tag. Lissamine imparts a positive charge on the morpholino to facilitate electroporation into retinal cells<sup>59</sup> and provides a fluorescent label to evaluate the effectiveness of the electroporation. Morpholino oligonucleotides were reconstituted to a concentration of 3 mM and 0.5  $\mu$ L MO solution was injected into the vitreous chamber of the left eye of each fish. Morpholino oligonucleotide injections were performed at 2 dpl, just before the onset of *neurod* expression in dividing progenitors.<sup>39</sup> Immediately after MO injection, eyes were electroporated across the dorsal-ventral axis following previously described protocols.<sup>60,61</sup> Briefly, electroporations each consisted of two pulses at 75 V for 50 ms and 1 second between pulses. To evaluate electroporation effectiveness, 10- $\mu$ m-thick cross-sections were cut through the central retina from each eye, slides were rinsed 3  $\times$  5 minutes with PBS, counterstained with 4',6-diamidino-2-phenylindole (DAPI), coverslipped and imaged immediately using epifluorescence to visualize lissamine distribution. Eyes with little or no fluorescent lissamine visible in retinal cells or that were damaged by the electroporation were discarded and not subjected to further analysis.

### Western Blot Analysis

Protein samples were obtained by pooling five adult retinas in lysis buffer containing protease and phosphatase inhibitors (5872; Cell Signaling Technology, Danvers, MA, USA). Proteins were separated in a 12% SDS-PAGE gel and transferred to a nitrocellulose membrane (Sigma-Aldrich Corp.). The membrane was blocked in 5% nonfat dry milk in PBS with 0.05% Tween-20 for 2 hours and then incubated overnight at 4°C with rabbit anti-NeuroD antibody<sup>7</sup> diluted 1:1000 in 2.5% blocking solution. Blots were rinsed with PBS+0.05% Tween-20 and incubated with horseradish peroxidase-conjugated secondary IgG (1:5000) for 1 hour. Bands were visualized using the enhanced chemiluminescence assay (ECL detection system;

Amersham Biosciences, Arlington Heights, IL, USA). As loading controls, blots were incubated in stripping buffer (21059; ThermoFisher Scientific) for 5 minutes, then blocked and labeled as described above with anti- $\beta$ -actin (1:5000, Calbiochem). Images were captured using the FluorChem E Imaging System (Bio-Techne, Minneapolis, MN, USA).

### Cell Cycle Analysis and Notch Inhibition in Adults

To label cells in S-phase of the cell cycle, at 3 dpl adult fish were injected intraperitoneally (IP) with 4 mM EdU to label proliferating cells. Animals then were returned to normal intensity and cyclical lighting, and were treated by swimming the fish for 24 hours in 5 mM BrdU before being killed at 6 or 7 dpl. To inhibit Notch signaling in adults, DAPT was diluted in aquarium water to 40  $\mu$ M (in 0.16% DMSO)<sup>37,62</sup> and fish were treated with DAPT or DMSO only from 4 to 6 dpl.

To analyze cell cycle exit, for each adult fish, BrdU, EdU, and BrdU/EdU colabeled cells were counted as described above and averaged from 3 nonadjacent sections from a single retina ( $n = 5-6$  fish per treatment). To quantify regenerated photoreceptors in adults, 10- $\mu$ m-thick sections were processed for in situ hybridization for *rhodopsin* (rods) or *pde6c* (cones) and labeled cells were counted manually. For both techniques in adult retinas, labeled cells were counted within a 500- $\mu$ m length of retina centered between the dorsal margin and optic nerve. Statistical significance was determined using a Student's *t*-test.

## RESULTS

### NeuroD Function Is Non-Cell-Autonomous

As a first test of the mechanisms by which NeuroD governs photoreceptor genesis and maturation, NeuroD was knocked down by injecting ATG-targeted MO and then cells were transplanted reciprocally between morphant and WT embryos. Dextran-labeled donor cells were analyzed for the presence or absence of EdU and markers of mature photoreceptors in the outer nuclear layer (ONL) of host retinas between 72 and 75 hpf, a time point at which photoreceptor genesis and maturation in the central retina are complete in the WT retina<sup>63</sup> but are incomplete following NeuroD knockdown.<sup>7</sup> To identify unambiguously mature rod photoreceptors derived from donor embryos, cells were transplanted from the Tg(XOPS:GFP) line, which expresses GFP under the control of the *Xenopus* rhodopsin promoter (Fadool 2003). To identify mature cone photoreceptors, the Zpr-1 antibody was used to label red and green double cones.<sup>64</sup> As a control, WT cells

TABLE 2. Fate of Donor Cells in NeuroD Knockdown/WT Chimeras

	WT Donor, Morphant Host	Morphant Donor, WT Host
% Dextran/EdU	15.3	0
Dextran/Zpr-1 (cones)	-	+
Dextran/GFP (rods)	-	+

Dextran-labeled WT donor cells transplanted into NeuroD knockdown (morphant) hosts remain in the cell cycle (15.3% were EdU-positive), and none differentiate into photoreceptors (-). In the reciprocal experiment, NeuroD morphant donor cells transplanted into WT embryos uniformly exit the cell cycle (all EdU-negative) and differentiate into photoreceptors (+).

were transplanted into WT hosts, which demonstrated that transplantation had no effect on cell cycle exit or photoreceptor maturation among donor-derived or host cells (Supplementary Fig. S1).

When WT cells were transplanted into NeuroD knockdown recipients, 15.3% of donor cells within the ONL were EdU-positive, and none of the donor cells expressed markers of mature rod or cone photoreceptors (Table 2; Figs. 1A, 1C, arrowheads). In the reciprocal experiment, when cells from NeuroD knockdown were transplanted into WT hosts, 0% of the donor cells contained EdU, and all donor cells within the ONL expressed markers of mature rod or cone photoreceptors (Table 2; Figs. 1B, 1D, arrowheads). These experiments demonstrated that cell cycle exit and maturation of transplanted cells are determined exclusively by the host retina, and, therefore, in cells of the photoreceptor lineages, NeuroD function is non-cell-autonomous. This indicates that NeuroD governs cell cycle exit and photoreceptor maturation via mechanisms mediated through cell-cell signaling.

### NeuroD Regulates Expression of Notch Signaling Molecules and Targets of the Notch Pathway

The data demonstrating that NeuroD function is non-cell-autonomous prompted us to investigate intercellular Notch signaling as a mediator of NeuroD function. Two time points were selected for analysis in the embryonic retina: 48 and 70 hpf. The 48-hpf time point allowed analysis of events leading up to the onset of photoreceptor genesis, which, outside the precocious ventral patch,<sup>65</sup> begins at approximately 48 hpf. The 70-hpf time point allowed analysis of events occurring between 48 and 70 hpf, corresponding to the onset and completion of photoreceptor genesis and maturation (see prior report<sup>63</sup>). In situ hybridization showed that in control retinas, the Notch receptor *notch1a*, the Notch ligands *deltaA* (*dla*) and *deltaD* (*dld*), and the Notch targets *ascl1a* and *ber4* are expressed throughout the central retina at 30 hpf (data not shown), but are downregulated in the central retina by 48 hpf (Fig. 2A, arrowheads) and restricted to the ciliary marginal zone (CMZ) by 70 hpf (Fig. 2B, arrowheads). However, NeuroD knockdown results in the failure of cells to downregulate these genes (Figs. 2A, 2B, arrowheads), and at 70 hpf this is most apparent in the developing ONL adjacent to the dorsal CMZ (Fig. 2B, circled region) where photoreceptor genesis is occurring. Corresponding qRT-PCR data from embryo heads show that at 48 and/or 70 hpf, NeuroD knockdown results in significantly increased expression of *notch1a*, *dla*, and *dld*, as well as several downstream targets of the Notch pathway including *cyclinD1*, *ascl1a*, *cyclinB1*, and *hes6*. The overall level of *ber4* expression is not significantly affected by NeuroD knockdown (Fig. 2C), and *ber4* expression appears to be





FIGURE 1. NeuroD function is non-cell-autonomous. Reciprocal transplant chimeric analysis showing the fates of dextran-labeled donor cells on the ONL. (A) Wild-type donor cells in NeuroD knockdown hosts remain in the cell cycle (EdU-positive) and do not differentiate into rod photoreceptors (arrowhead; Tg[XOPS:GFP]-negative). (B) NeuroD knockdown Tg(XOPS:GFP) donor cells in WT hosts exit the cell cycle (EdU-negative) and differentiate into rods (arrowhead; Tg[XOPS:GFP]-positive). (C) Wild type donor cells in NeuroD knockdown hosts remain in the cell cycle (EdU-positive) and do not differentiate into cones (arrowhead; Zpr-1-negative). (D) NeuroD knockdown donor cells in WT hosts exit the cell cycle (EdU-negative) and differentiate into cones (arrowhead; Zpr-1-positive). GCL, ganglion cell layer. Scale bar: 50  $\mu$ m.

unaffected in the brain (data not shown). This suggests that NeuroD regulation of *ber4* is restricted primarily to the retina and suggests the possibility that *ber4* may be regulated by other mechanisms in addition to the Notch pathway. These data demonstrated that in the developing retina, NeuroD negatively regulates intercellular Notch signaling, thereby negatively regulating downstream targets of the Notch pathway.

### NeuroD is Not Required for Photoreceptor Fate Specification or Progenitor Survival

Two questions remain unanswered from previous studies: Is NeuroD required for photoreceptor cell fate specification, and does the small eye phenotype following NeuroD knockdown result from increased apoptosis? To answer the first question, in situ hybridization was used to determine if NeuroD knockdown alters the expression of the canonical markers of the photoreceptor lineages *crx* and *nr2e3*. In the developing retina, *crx* and *nr2e3* are expressed in mitotic photoreceptor progenitors, then *nr2e3* is downregulated in postmitotic cones.<sup>66,67</sup> NeuroD knockdown does not prevent the expression of these genes in either dividing or postmitotic cells (Fig. 3) and, due to the lack of mature cones following NeuroD knockdown, results in *nr2e3* expression in a higher number of cells (Fig. 3G).




**FIGURE 2.** NeuroD negatively regulates Notch signaling molecules and downstream targets of the Notch pathway. **(A, B)** In situ hybridizations showing the normal retinal expression of *neurod* compared to *notch1a*, *dla*, *dld*, *ascl1a*, and *her4* at 48 **(A)** and 70 **(B)** hpf in control embryos and following NeuroD knockdown. *Arrowheads* indicate zones of expression; *circled regions* highlight expression differences at 70 hpf in and around the developing ONL and INL adjacent to the dorsal CMZ. *Scale bars*: 50  $\mu$ m. **(C)** Quantitative RT-PCR data indicating the fold differences in expression between control embryos (*black horizontal line*) and embryos following NeuroD knockdown at 48 and 70 hpf. \* $P < 0.05$ , Student's *t*-test. *Error bars*: standard deviation.

These observations indicate that NeuroD is not required for photoreceptor fate specification. To address the second issue, TUNEL labeling revealed that at 70 hpf, the number of apoptotic cells in the retina does not change following NeuroD knockdown (Supplementary Fig. S2). This indicates that during the temporal window of photoreceptor genesis, NeuroD is not required for the survival of photoreceptor progenitors. This suggests that the reduced eye diameter at 70 hpf resulting from NeuroD knockdown likely is caused by

the lack of photoreceptor differentiation rather than cell death.

### CRISPR/Cas9 Targeted Mutation of *neurod* Phenocopies Morpholino Knockdown and Identically Affects Notch Signaling

The zebrafish community has raised concerns about potential off-target effects of MOs and cautioned that MOs should be




**FIGURE 3.** During embryonic development, NeuroD knockdown does not affect expression of the photoreceptor lineage markers *nr2e3* and *crx*. In control retinas at 72 hpf, *crx* (A) and *nr2e3* (E) are expressed in postmitotic photoreceptors (black arrowheads) and in EdU+ progenitors (white arrowheads) adjacent to the CMZ that have not yet exited the cell cycle. Likewise, following NeuroD knockdown (*neuroD* MO), *crx* (B–D), and *nr2e3* (F–H) are expressed in postmitotic photoreceptors (black arrowheads) and in EdU+ progenitors (white arrowheads) distributed throughout the ONL. Scale bar: 50  $\mu$ m.

used only if targeted gene mutation produces the same phenotype.<sup>68,69</sup> To determine if targeted mutation of *neuroD* results in the same retinal phenotype as NeuroD knockdown, preliminary experiments were undertaken using CRISPR/Cas9 genome editing to create targeted mutations in the *neuroD* gene. To characterize mutations in individual embryos displaying a retinal phenotype, DNA was collected at 70 hpf, PCR amplified across the sgRNA target site and subcloned into pGEM-T Easy (Promega). Six to nine clones were sequenced per embryo, and between one and three indels were identified in each embryo from distinct clones. At least one WT DNA strand was detected in each embryo, indicating mosaicism at the targeted sequences. Mutations ranged from deletions of 1 to 69 nucleotides and/or insertions of 1 to 2 nucleotides, all starting or ending within the center of the sgRNA target sequence. Most indels introduced putative premature stop codons. CRISPR/Cas9-injected embryos that tested negative for insertions/deletions (indels) had normal retinas, identical to WT uninjected embryos (Fig. 4A). Of 14 injected embryos analyzed, 12 (86%) tested positive for indels, and 5 of these (42%) had a retinal phenotype consistent with that observed following NeuroD knockdown using MOs. This phenotype included reduced eye size (Fig. 4A), excess BrdU+ cells in the ONL and paucity of mature photoreceptors (Fig. 4A, cf. Figs. 1A, 1C; arrowheads). NeuroD knockdown and targeted mutation of *neuroD* also produced identical changes to expression of genes in the Notch pathway. At 70 hpf, cells in embryos with *neuroD* indels had increased expression of *notch1a*, *dla*, *dld*, *ascl1a*, and *ber4* in the central retina (Fig. 4B, arrowheads and circled regions). The phenotypes combined with the sequencing results suggest that embryos with *neuroD* indels are equivalent to embryos in which NeuroD has been knocked down. These data demonstrated that reductions in NeuroD caused by targeted mutation or morpholino

knockdown cause identical retinal phenotypes and increased expression of Notch pathway genes.

### Notch Signaling Governs NeuroD-Mediated Cell Cycle Exit, But Not Photoreceptor Maturation

As a test of the hypothesis that Notch signaling mediates NeuroD function, Notch signaling was inhibited by exposing embryos to the gamma secretase inhibitor, DAPT, between 48 and 70 hpf following NeuroD knockdown. DAPT treatment significantly reduced expression of the Notch targets *ber4* and *ascl1a* (Fig. 5A), restoring an expression pattern similar to controls (Fig. 5B, cf. Figs. 2A–C; arrowheads). These results confirmed that DAPT treatment inhibited the Notch pathway, and are interpreted to show that NeuroD regulation of *ber4* and *ascl1a* is mediated through Notch signaling. Similarly, following NeuroD knockdown, DAPT treatment reduced the number of BrdU-positive progenitors to control levels (Figs. 5C, 5D), demonstrating that in the absence of NeuroD, inhibition of Notch signaling is sufficient to drive photoreceptor progenitors out of the cell cycle. However, DAPT treatment was not sufficient to rescue photoreceptor maturation (Fig. 5D). To determine if the activation of Notch signaling in the presence of the normal expression of NeuroD is sufficient to prevent cycle exit and photoreceptor maturation, Tg(hsp70:GGFF)  $\times$  Tg(UAS:*myc-notch1a:intra*) embryos were heat-shocked at 52 to 53 hpf to conditionally activate the Notch pathway. Notch activation (Notch gain-of-function [GOF]) caused substantial upregulation of the direct Notch target gene *ber4* throughout the lens and retina (Fig. 6A, arrowheads), confirming strong upregulation of Notch pathway activity. Compared to controls, Notch GOF caused modest increases in the expression of *cyclinD1* and *ascl1a* in the central retina (Fig. 6A, arrowheads), resulted in significantly more EdU+ progenitors in the ONL, and caused a marked




**FIGURE 4.** CRISPR/Cas9 targeted mutation of *neuroD* phenocopies NeuroD knockdown and similarly affects Notch signaling. (A) EdU and Zpr-1 (cones) labeling in retinas of uninjected and WT injected (T7 endonuclease [EN]-negative) embryos, compared to retinas of 3 embryos positive for insertions and/or deletions in the *neuroD* gene (T7 endonuclease [EN]-positive). Zpr-3 labeling (rods) was affected identically (data not shown). As a relative comparison of photoreceptor progenitors still in the cell cycle in the central retina, *arrowheads* indicate the central-most EdU+ cells in the ONL. (B) In situ hybridization showing retinal expression of *notch1a*, *dla*, *dld*, *ascl1a*, and *her4* in embryos with *neuroD* indels compared to uninjected controls. *Arrowheads* indicate zones of expression; *circled regions* highlight expression differences in and around the developing ONL and INL adjacent to the dorsal CMZ. *Scale bars*: 50  $\mu$ m.

decrease in photoreceptor maturation (Figs. 6B, 6C). Furthermore, identical to NeuroD knockdown, Notch GOF resulted in reduced eye size (Figs. 6A, 6B) and no increased apoptosis (data not shown). These results demonstrated that conditional

activation of the Notch signaling pathway is sufficient to prevent photoreceptor genesis, recapitulating NeuroD knockdown. Together, the Notch inhibition and GOF data are interpreted to show that cell cycle exit among photoreceptor






**FIGURE 5.** Inhibition of Notch signaling reduces *ber4*, *ascl1a*, and ONL cell proliferation following NeuroD knockdown, but does not rescue photoreceptor differentiation. (A) Quantitative RT-PCR data showing fold changes in *ber4* and *ascl1a* expression following NeuroD knockdown in embryos treated 48 to 70 hpf with DAPT, compared to control embryos treated with DMSO only (solid black line). (B) In situ hybridization showing retinal expression of *ber4* and *ascl1a* following NeuroD knockdown in embryos treated 48 to 70 hpf with DMSO only compared to embryos treated with DAPT. Dashed lines and arrowheads indicate zones of expression. (C) Percentage of retinal cells labeled with BrdU following NeuroD knockdown in embryos treated with DMSO only, compared to embryos treated with DAPT, and with embryos injected with MM control morpholinos treated with DMSO only. (D) BrdU and Zpr-1 (cones) labeling following NeuroD knockdown in embryos treated with DMSO only compared to embryos treated with DAPT; note the absence of cones in both images. Scale bars: 50  $\mu$ m. Actual *P* values shown on graphs; \*\**P* < 0.005, \*\*\**P* < 0.0005, Student's *t*-test. Error bars: standard deviation.

progenitors is governed through repression of Notch signaling by NeuroD, but that NeuroD governs photoreceptor maturation through mechanisms independent of the Notch pathway.

### Time Course of *neurod* Expression in the Adult Regenerating Retina

As a first step to understand NeuroD function during photoreceptor regeneration, following photoreceptor ablation with bright light, *neurod* expression was characterized in relation to *pax6a*, a marker of dividing progenitors as well as

mature amacrine and ganglion cells, and the photoreceptor lineage markers *crx* and *nr2e3*. Within 24 to 48 hours after light onset (hpl), a subset of Müller glia divides once, each producing a mitotic daughter cell that subsequently divides multiple times to produce a cluster of neurogenic progenitors.<sup>50</sup> Therefore, in the present study, 48 hpl was the earliest time point analyzed for progenitor markers. In situ hybridization and immunohistochemistry showed that the progenitor cell marker *pax6a* and the photoreceptor lineage marker *nr2e3* are expressed in proliferating cell nuclear antigen-positive (PCNA+) cells by 56 hpl (Fig. 7). In comparison,




**FIGURE 6.** Conditional activation of the Notch pathway up-regulates Notch target expression and prevents cell cycle exit and photoreceptor genesis. **(A)** In situ hybridization showing retinal expression of the Notch targets *her4*, *cyclinD1*, and *ascl1a* in control embryos compared to Notch GOF embryos heat-shocked 52-53 hpf. **(B)** EdU, Zpr-1, and Myc labeling in 70 hpf control (top panels: Tg[hsp:GGFF] only) and Notch GOF (Tg[hsp:GGFF] × Tg[UAS:*myc-notch1a:intra*]) embryos heat-shocked 52-53 hpf. **(C)** Quantification of EdU+ cells in the ONL of 70 hpf control embryos ( $n = 5$ ) compared to Notch GOF embryos ( $n = 8$ ) heat-shocked 52 to 53 hpf. Scale bars: 50  $\mu$ m. \*\*\* $P = 0.00003$ , Student's  $t$ -test. Error bars: standard deviation.

*neurod* mRNA was first detectable in a subset of PCNA+ progenitors at 68 hpl (Fig. 7), slightly earlier than the 72 hpl time point identified in earlier studies.<sup>39</sup> The photoreceptor lineage marker *crx* is constitutively expressed throughout the inner nuclear layer (INL) and ONL, and also was first noticeably upregulated in a subset of PCNA+ progenitors only after 68 hpl (Fig. 7). NeuroD knockdown did not prevent *nr2e3* or *crx* expression (Supplementary Fig. S3), indicating that during

photoreceptor regeneration, NeuroD is not required for photoreceptor fate specification among the Müller glia-derived progenitors.

At 72 hpl and later, the Tg(*neurod:egfp*) line is an accurate reporter of endogenous *neurod* expression<sup>39</sup> and showed that at 72 hpl, all BrdU+/EGFP+ (*neurod*+) photoreceptor progenitors express *pax6a*, *nr2e3*, and *crx* (Supplementary Fig. S4). However, by 7 dpl, when regenerated photoreceptors are




**FIGURE 7.** Earliest detectable expression of *pax6*, *nr2e3*, *neurod*, and *crx* in Müller glia-derived progenitors. **(A)** *pax6* and *nr2e3* expression is detectable in PCNA+ progenitors as early as 56 hpl (arrowheads). **(B)** *neurod* expression is detectable in PCNA+ cells as early as 68 hpl, but in Tg(*neurod:egfp*) fish, EGFP immunolabeling is not yet detectable in these cells (black arrowheads); *crx* is constitutively expressed throughout the INL and at 68 hpl is first detectable at higher levels in PCNA+ progenitors (white arrowheads). Scale bars: 50  $\mu$ m.

exiting the cell cycle and differentiating,<sup>29,60</sup> *pax6a* is downregulated in all photoreceptors while *neurod* and *nr2e3* are downregulated in newly postmitotic cones but continue to be expressed in rods (Supplementary Fig. S4). This pattern persists in the fully regenerated retina at 14 dpl (Supplementary Fig. S5) and suggests a role for NeuroD during maturation and/or maintenance of rods.

### Following Photoreceptor Ablation, NeuroD Governs Cell Cycle Exit and Photoreceptor Regeneration

The knowledge that NeuroD governs cell cycle exit among photoreceptor progenitors in the embryonic retina prompted us to determine if NeuroD functions similarly during photoreceptor regeneration. To test this, MOs were used to knock down NeuroD protein and labeling with two different cell cycle markers, EdU and BrdU, was used to analyze cell cycle exit. Morpholinos were injected at 48 hpl, before the onset of *neurod* expression, to prevent translation of NeuroD protein. Western blots confirmed efficient knockdown of NeuroD protein at 4 dpl (14.5% of control), but recovery of NeuroD protein by 7 dpl (107% of control; Fig. 8A). EdU was

administered systemically at 3 dpl, when photoreceptor progenitors are proliferating in the INL,<sup>29</sup> and fish were exposed to BrdU from 5 to 6 dpl or from 6 to 7 dpl, when photoreceptor progenitors are in the nascent ONL and normally exiting the cell cycle.<sup>29,60</sup> This paradigm differentially labels cells in S-phase of the cell cycle at each time point and the percentage of cells colabeled with EdU and BrdU was used as a proxy for the percentage of cells that remained in the cell cycle between 3 and 6 dpl. Compared to controls at 6 and 7 dpl, NeuroD knockdown resulted in a higher percentage of proliferating cells colabeled with EdU and BrdU (Figs. 8B, 8C). There was a greater difference between control and knock-down retinas at 6 than at 7 dpl (Fig. 8B), perhaps reflecting the recovery of NeuroD by 7 dpl (cf. Fig. 8A). This demonstrates that NeuroD knockdown prevents photoreceptor progenitors from exiting the cell cycle and is interpreted to show that in the injured, regenerating retina, NeuroD governs cell cycle exit among the Müller glia-derived photoreceptor progenitors. To determine if the results described above impact photoreceptor regeneration, *in situ* hybridization was used at 6 and 7 dpl to label regenerated cones (*pde6c*) and rods (*rhodopsin*). At 6 dpl, NeuroD knockdown resulted in fewer regenerated cones and rods (Figs. 9A, 9B); however, by 7 dpl, cone regeneration



**FIGURE 8.** NeuroD is required for photoreceptor progenitors to exit the cell cycle. (A) Following photoreceptor ablation and morpholino injection/electroporation at 2 dpl, Western blot indicates strong knockdown of NeuroD protein (*neuroD* MO) at 4 dpl compared to control MO or uninjected retinas, but recovery of NeuroD protein by 7 dpl. (B) When fish are injected with EdU at 3 dpl and exposed to BrdU from 5 to 6 or from 6 to 7 dpl, NeuroD knockdown results in a greater percentage of progenitors that are colabeled with EdU and BrdU at 6 and 7 dpl, indicating that fewer cells have exited the cell cycle; Student's *t*-test, \* $P < 0.05$ . (C) Corresponding images show the increase in EdU+/BrdU+ colabeled cells (arrowheads) following NeuroD knockdown. Scale bar: 50  $\mu$ m.


recovers along with the recovery of NeuroD (Figs. 9A, 9C). These data demonstrated that, following NeuroD knockdown, the failure of progenitors to exit the cell cycle is sufficient to prevent or delay photoreceptor regeneration.

### NeuroD Governs Photoreceptor Regeneration Through Delta-Notch Signaling

The data showing that NeuroD governs cell cycle exit through Notch signaling in the developing retina led us to investigate Notch signaling as a putative mediator of NeuroD function during photoreceptor regeneration. First, in situ hybridization was used to characterize the timing of Notch pathway gene expression in photoreceptor progenitors. At 72 hpl, *deltaA*, *notch1a*, and *her4* are coexpressed in BrdU+ cells (Figs. 10A, 10B). By 7 dpl, most progenitors that were labeled with BrdU at 3 dpl have exited the cell cycle (cf. Fig. 8) and no longer express Notch pathway genes (Fig. 10C). These observations showed that within photoreceptor progenitors, Notch path-

way gene expression coincides with proliferation and is downregulated during photoreceptor differentiation.


Downregulation of Notch pathway genes, concomitant with increasing *neurod* expression and cell cycle exit, suggests that Notch signaling and NeuroD may affect the cell cycle in opposing ways. To determine if Notch signaling genes are regulated downstream of NeuroD, in situ hybridization was used to compare expression of the Notch signaling molecules *notch1a*, *deltaA*, and *deltaD*, and the Notch targets *her4* and *ascl1a*, in control and experimental retinas. At 4 dpl, a time point when photoreceptor progenitors are proliferating and few have exited the cell cycle, NeuroD knockdown has no discernable effect on Notch pathway gene expression (data not shown). However, at 6 dpl, when regenerated photoreceptors are normally differentiating (cf. Figs. 8, 9), NeuroD knockdown results in increased expression of Notch signaling genes and Notch targets (Fig. 11). This indicates that NeuroD knockdown prevents the normal downregulation of Delta-Notch signaling



**FIGURE 9.** NeuroD is required for photoreceptor regeneration. (A) Compared to controls (*solid black line*), NeuroD knockdown results in reduced rod and cone regeneration by 6 dpl, but only reduced rod regeneration by 7 dpl. Student's *t*-test,  $*P < 0.05$ . Corresponding images at 6 dpl (B) and 7 dpl (C) show in situ hybridizations for cones (*pde6c*) and rods (*rhodopsin*). Scale bars: 50  $\mu$ m.

among progenitors and that during photoreceptor regeneration NeuroD negatively regulates Delta-Notch signaling. To determine if Notch signaling mediates NeuroD function, following photoreceptor ablation and NeuroD knockdown,

fish were treated with the gamma secretase inhibitor, DAPT, from 4 to 6 dpl. Proliferating cells were labeled with EdU (IP injection) at 3 dpl and with BrdU (immersion) at 5 to 6 dpl. Colabeling analysis was performed as described above to




**FIGURE 10.** Notch signaling molecules are expressed in proliferating photoreceptor progenitors. (A) In situ hybridizations show that at 72 hpl, the Notch ligand *deltaA*, the Notch receptor *notch1a*, and the direct Notch target *her4* are expressed in proliferating (BrdU+) photoreceptor progenitors (arrowheads). (B) Double in situ hybridization shows colocalization of *deltaA* and *notch1a* in the same cells (arrowheads). (C) By 7 dpl, *deltaA*, *notch1a*, and *her4* are no longer expressed in the cells that were proliferating at 3 dpl (labeled with BrdU). Scale bars: 50  $\mu$ m.

determine the proportion of progenitors that remained in the cell cycle. DAPT treatment reduced expression of *her4* and *ascl1a*, confirming that Notch signaling was inhibited (Supplementary Fig. S6), reduced the percentage of progenitors colabeled with EdU and BrdU to control levels (Figs. 12A, 12B) and fully rescued the deficiency in photoreceptor regeneration induced by NeuroD knockdown (Figs. 12C, 12D). These results indicate that inhibiting Notch signaling is sufficient to rescue NeuroD knockdown, and that NeuroD governs photoreceptor regeneration through negatively regulating Notch signaling.

## DISCUSSION


Photoreceptor genesis in the vertebrate retina requires precise regulation of events that determine progenitor fate, regulate the timing of cell cycle exit, and induce differentiation and maturation.<sup>1,70-73</sup> During retinal development in zebrafish the bHLH transcription factor NeuroD is expressed in mitotic photoreceptor progenitors and nascent photoreceptors,<sup>13,14,39</sup> and within the photoreceptor lineage, NeuroD governs cell cycle exit and photoreceptor maturation.<sup>7</sup> Photoreceptor genesis normally is complete by 70 hpf<sup>63</sup>; however, in the absence of NeuroD, even though cellular laminae form,



**FIGURE 11.** NeuroD negatively regulates Notch signaling molecules and targets of the Notch pathway. At 6 dpl in both the INL and ONL, in situ hybridizations indicate that NeuroD knockdown results in increased expression of the Notch receptor *notch1a*, the Notch ligands *deltaA* and *deltaD*, and the Notch pathway targets *her4* and *ascl1a*. Scale bar: 50  $\mu$ m.

photoreceptor progenitors fail to exit the cell cycle and mature photoreceptors are absent.<sup>7</sup> Following conditional ectopic expression of NeuroD, nearly all dividing cells are forced out of the cell cycle, *cyclinD1*, *cyclinB1*, and *cyclinE1* are downregulated and cyclin kinase inhibitors, *p27* and *p57*, are upregulated.<sup>7</sup> In the adult retina following photoreceptor ablation, *neurod* is expressed in Müller glia-derived mitotic

progenitors.<sup>39</sup> This suggests that NeuroD also has a role in photoreceptor regeneration, but this has not been investigated. The goal of the current study was to identify the molecular mechanisms through which NeuroD functions and to mechanistically link NeuroD to the mechanisms that govern the cell cycle during photoreceptor genesis in embryos and photoreceptor regeneration in adults.



**FIGURE 12.** NeuroD governs cell cycle exit and photoreceptor regeneration through Notch signaling. **(A)** Compared to controls (control MO treated with DMSO vehicle), NeuroD knockdown (*neurod* MO treated with DMSO vehicle) results in an increased percentage of EdU+/BrdU+ colabeled cells, but inhibition of Notch signaling with the gamma secretase inhibitor DAPT from 4 to 6 dpl (*neurod* MO, DAPT) restores the percentage of EdU+/BrdU+ colocalized cells to control levels. Student's *t*-test, \* $P < 0.05$ . **(B)** Corresponding images at 6 dpl show the reduction in EdU+/BrdU+ colabeled cells (arrowheads) in *neurod* morphants following DAPT treatment. **(C)** NeuroD knockdown results in fewer regenerated cone and rod photoreceptors, but inhibition of Notch signaling with the gamma secretase inhibitor DAPT from 4 to 6 dpl (*neurod* MO, DAPT) restores the number of regenerated photoreceptors to control levels. Student's *t*-test, \* $P < 0.05$ . **(D)** Corresponding images at 6 dpl show in situ hybridizations for cones (*pde6c*) and rods (*rhodopsin*). Scale bars: 50  $\mu$ m.

In the embryonic retina, genetic mosaic analysis showed that NeuroD regulates cell cycle exit and photoreceptor maturation through non-cell-autonomous mechanisms. This indicates that NeuroD governs photoreceptor genesis and maturation through the regulation of intercellular signaling events. Transcription factors by their nature function cell-autonomously by binding DNA to regulate gene transcription, but they also can function non-cell-autonomously by regulating intercellular signaling molecules.<sup>29,36,74–78</sup> This also is true for bHLH factors that regulate intercellular signaling pathways in the retina.<sup>23,29,36,76</sup> There also are rare examples of transcription factors functioning as secreted molecules.<sup>79–81</sup> In vitro, naked NeuroD protein can be transported through cell membranes and activate transcriptional events.<sup>82</sup> However, there is no evidence in vivo for the secretion of NeuroD or any other bHLH transcription factor, and we infer that NeuroD is not secreted in the developing or regenerating zebrafish retina.

Notch signaling was investigated as the putative non-cell-autonomous mediator of NeuroD function and our data were interpreted to show that NeuroD governs cell cycle exit in photoreceptor progenitors through repression of Delta-Notch signaling. Gamma-secretase inhibitors, such as DAPT, effective-

ly inhibit Notch pathway activation in the zebrafish retina and other models<sup>18,21,34,37,83</sup>; however, it should be noted that gamma-secretase is necessary for other signaling mechanisms, such as through the amyloid precursor protein (APP).<sup>83</sup> Therefore, it is possible that some of the effects of DAPT treatment, such as regional specific expression patterns of *ber4* and *ascl1a*, result from interference with other pathways in addition to Notch signaling. However, our Notch GOF data in the embryo reciprocate the DAPT data with regard to cell cycle exit and photoreceptor genesis and, therefore, strongly support our conclusions. Our findings are consistent with the conserved role of Notch signaling to prevent differentiation and maintain progenitors in a proliferative state.<sup>18–21,84,85</sup> Several studies implicate *cyclinD1*, *cyclinD3*, and *cdk* genes as direct regulatory targets of the Notch signaling pathway, whereby Notch signaling promotes progression from G1 to S phases of the cell cycle.<sup>86–89</sup> Notch signaling also indirectly regulates *cyclinB1* expression.<sup>90</sup> Therefore, transcriptional repression of Notch signaling by NeuroD provides a mechanistic link between NeuroD and cell cycle regulation through *cyclinB1* and *cyclinD1* (see also the report by Ochocinska and Hitchcock<sup>7</sup>).



Earlier work determined that NeuroD governs photoreceptor genesis and maturation in the developing zebrafish retina.<sup>7</sup> However, that study did not determine if NeuroD is required for specification of the photoreceptor lineage or for cell survival. The canonical photoreceptor lineage markers *nr2e3* and *crx* are expressed in mitotic photoreceptor progenitors.<sup>66,67,91–93</sup> NeuroD knockdown does not prevent expression of either *crx* or *nr2e3*, indicating that NeuroD is not required to specify the photoreceptor lineage. At 70 hpf, NeuroD knockdown also did not alter the number of apoptotic cells in the developing retina, indicating that during the time of normal photoreceptor genesis, NeuroD is not required for survival of photoreceptor progenitors. Our transient knockdown of NeuroD, however, cannot determine if NeuroD is required for survival of photoreceptors or progenitors beyond this time point. In the mouse retina, NeuroD1 is not required for photoreceptor lineage specification, progenitor survival, or photoreceptor genesis, but is required for opsin selection among cones<sup>11</sup> and for survival of a subset of mature rod photoreceptors.<sup>10</sup> In the mouse hippocampus and olfactory bulb, NeuroD1 is required for survival and maturation of adult-born neurons.<sup>94</sup> These observations suggest that NeuroD also might have a role in neuron survival in zebrafish and will be a focus of future work.

The zebrafish community has raised concerns regarding the use of MOs as experimental tools due to the possibility of unidentifiable off-target effects.<sup>69</sup> Some have shown that in many cases gene mutation and morpholino-based protein knockdown do not produce the same developmental phenotype.<sup>68</sup> These studies conclude, however, that when used appropriately and when morpholino-based knockdown and genetic mutation produce the same phenotype, morpholinos can be effective tools for research. In the present study, morpholino-based knockdown of NeuroD and CRISPR/Cas9-based mutation of *neurod* produce the same retinal phenotype. Despite the expected cellular mosaicism of the gene disruption in CRISPR/Cas9 injected embryos, nearly half of these embryos have a retinal phenotype consistent with NeuroD knockdown. Both approaches show that NeuroD is required for photoreceptor progenitors to exit the cell cycle and photoreceptors to differentiate, and that NeuroD regulates Notch pathway gene expression. This validates the results from previous studies that investigated NeuroD function,<sup>7</sup> provides two independent methods to test NeuroD function and validates the use of morpholinos in the present study.

Among photoreceptor progenitors, NeuroD links cell cycle exit, differentiation, and maturation, but our data indicated that NeuroD regulates these events through separate non-cell-autonomous mechanisms. In the embryo, inhibition of Notch signaling following NeuroD knockdown is sufficient to rescue defects in cell cycle exit, but is not sufficient to restore photoreceptor maturation. This observation differs from *Id2a* knockdown in which inhibition of Notch signaling rescues defects in cell cycle exit and photoreceptor maturation.<sup>21</sup> These disparate results can be explained by the normal retinal expression of *neurod* following *Id2a* knockdown (Taylor SM, Hitchcock PF, unpublished data, 2014), suggesting that, independent of Notch signaling, photoreceptor maturation is critically dependent on the normal expression of *neurod*. These data demonstrate that cell cycle exit and photoreceptor maturation are regulated by NeuroD, but are regulated through different pathways. Our findings are consistent with data showing that, as in the *ying/brg1* mutation in the developing retina,<sup>43,95</sup> fate specification, cell cycle withdrawal, and differentiation can be uncoupled.

Following photoreceptor ablation in the adult retina, *nr2e3* is expressed before *neurod* and NeuroD knockdown does not prevent either *nr2e3* or *crx* expression. These observations

indicated that in the regenerating retina, as in the embryo, NeuroD does not specify the photoreceptor fate. As regenerated photoreceptors differentiate, *neurod* and *nr2e3* are downregulated in cones but continue to be expressed in rods. This suggests a role for NeuroD in terminal differentiation and/or photoreceptor maintenance, independent of Notch signaling. The absence of *neurod* expression from mature cones has been documented in embryonic and adult retinas,<sup>14,39</sup> although recent data indicate rhythmic, circadian expression in mature cones peaking in the light phase and in mature rods peaking in the dark phase.<sup>96</sup> Our data were all collected during the light phase, indicating that loss of *neurod* from newly regenerated cones results from regenerative mechanisms rather than circadian rhythmicity.

Following photolytic lesioning, NeuroD knockdown prevents cell cycle exit and photoreceptor regeneration and results in increased expression of Notch pathway genes. These defects are rescued by Notch inhibition, and this is interpreted to show that, in the light-damaged retina, NeuroD governs cell cycle exit and photoreceptor regeneration through Notch signaling. This shows that Notch signaling prevents cell cycle exit in photoreceptor progenitors, contrasting with the effect of Notch signaling on Müller glia to maintain these cells in a nonproliferative, differentiated state.<sup>34</sup> This suggests that Notch signaling might regulate the single, asymmetrical division of Müller glia, causing the parent cell to exit the cell cycle while simultaneously causing the daughter cell to proliferate. This is consistent with the role of Notch in the zebrafish retina to promote gliogenesis in some progenitors while maintaining others in a proliferative state<sup>18,20</sup> and the requirement for Notch in the asymmetrical division of progenitors that give rise to glia/neuron sibling pairs in the *Drosophila* CNS.<sup>97</sup> Within 24 to 48 hours after injury, the time frame when Müller glia divide, the Notch pathway genes *notch1a*, *deltaA*, and *her4* are strongly upregulated in the retina and remain upregulated through 7 dpl.<sup>37</sup> We show that by 72 hpl, Notch pathway genes are expressed in all proliferating photoreceptor progenitors. This differs from observations after physical damage to all retinal layers where, 4 days after retinal injury, *notch1a* was expressed in all BrdU-positive cells while *delta* genes were “preferentially expressed” in apposing BrdU-negative cells.<sup>37</sup> The disparity between our study and that of Wan et al.<sup>37</sup> may be due to their approach of damaging all retinal layers, thus generating a more heterogeneous mix of progenitor types and variability in Notch pathway gene expression. We also showed that among photoreceptor progenitors in the developing and regenerating retina, NeuroD negatively regulates *ascl1a* through repression of Notch signaling. This indicates that in photoreceptor progenitors, Notch signaling positively regulates *ascl1a*, contrasting with the effect of Notch signaling to negatively regulate *ascl1a* in Müller glia.<sup>37</sup> This suggests that, in photoreceptor progenitors, *Ascl1a* functions downstream of Notch in the transition from proliferation to differentiation and suggests a mechanism through which Notch signaling may regulate the cell cycle differently in Müller glia and their progeny.

In conclusion, our data demonstrated a conserved role for NeuroD during photoreceptor development and regeneration in which Notch signaling is a mechanistic link between NeuroD and cell cycle exit. We also showed that during embryonic development, NeuroD functions non-cell-autonomously and governs photoreceptor maturation through pathways that do not involve Notch signaling. Future work will focus on identifying the mechanisms, during development and regeneration, that function upstream of NeuroD and through which NeuroD governs photoreceptor maturation.

## Acknowledgments

The authors thank Ann C. Morris, University of Kentucky, for providing Tg(UAS:*myc-notch1a:intra*) fish and for scientific consult; Laura Kakuk-Atkins, Dilip Pawar, and Zachary Stamlis for technical assistance; and Christine Crilly for administrative assistance.

Supported by grants from the National Institutes of Health/National Eye Institute ([NIH/NEI], Bethesda, MD, USA), R01 EY07060 (PFH), F32 EY023129 (SMT), T32 EY013934 (SMT), Core Grant P30EYO7003 (University of Michigan), Core Grant P30RY004068 (Wayne State University), NIH EY017753 (JMF), NIH EY020106 (KAD), grants from Fight for Sight and the Midwestern Eye-Bank (PFH), and Research to Prevent Blindness (Wayne State University). Fish lines and reagents provided by ZIRC were supported by NIH-NCRR Grant P40 RR01. The authors alone are responsible for the content and writing of the paper.

Disclosure: **S.M. Taylor**, None; **K. Alvarez-Delfin**, None; **C.J. Saade**, None; **J.L. Thomas**, None; **R. Thummel**, None; **J.M. Fadool**, None; **P.F. Hitchcock**, None

## References

- Bassett EA, Wallace VA. Cell fate determination in the vertebrate retina. *Trends Neurosci.* 2012;35:565-573.
- Brzezinski JA, Kim EJ, Johnson JE, Reh TA. Ascl1 expression defines a subpopulation of lineage-restricted progenitors in the mammalian retina. *Development.* 2011;138:3519-3531.
- Pollak J, Wilken MS, Ueki Y, et al. ASCL1 reprograms mouse Müller glia into neurogenic retinal progenitors. *Development.* 2013;140:2619-2631.
- Mao C-A, Cho J-H, Wang J, et al. Reprogramming amacrine and photoreceptor progenitors into retinal ganglion cells by replacing Neurod1 with Atoh7. *Development.* 2013;140:541-551.
- Akagi T, Inoue T, Miyoshi G, et al. Requirement of multiple basic helix-loop-helix genes for retinal neuronal subtype specification. *J Biol Chem.* 2004;279:28492-28498.
- Conte I, Marco-Ferreres R, Beccari L, et al. Proper differentiation of photoreceptors and amacrine cells depends on a regulatory loop between NeuroD and Six6. *Development.* 2010;137:2307-2317.
- Ochocinska MJ, Hitchcock PF. NeuroD regulates proliferation of photoreceptor progenitors in the retina of the zebrafish. *Mech Devel.* 2009;126:128-141.
- Lee JE, Hollenberg SM, Snider L, Turner DL, Lipnick N, Weintraub H. Conversion of *Xenopus* ectoderm into neurons by NeuroD, a basic helix-loop-helix protein. *Science.* 1995;268:836-844.
- Farah MH, Olson JM, Susic HB, Hume RI, Tapscott SJ, Turner DL. Generation of neurons by transient expression of neural bHLH proteins in mammalian cells. *Development.* 2000;127:693-702.
- Morrow EM, Furukawa T, Lee JE, Cepko CL. NeuroD regulates multiple functions in the developing neural retina in rodent. *Development.* 1999;126:23-36.
- Liu H, Etter P, Hayes S, et al. NeuroD1 regulates expression of thyroid hormone receptor 2 and cone opsins in the developing mouse retina. *J Neurosci.* 2008;28:749-756.
- Yan R-T, Wang S-Z. Requirement of neuroD for photoreceptor formation in the chick retina. *Invest Ophthalmol Vis Sci.* 2004;45:48-58.
- Hitchcock P, Kakuk-Atkins L. The basic helix-loop-helix transcription factor neuroD is expressed in the rod lineage of the teleost retina. *J Comp Neurol.* 2004;477:108-117.
- Ochocinska MJ, Hitchcock PF. Dynamic expression of the basic helix-loop-helix transcription factor neuroD in the rod and cone photoreceptor lineages in the retina of the embryonic and larval zebrafish. *J Comp Neurol.* 2007;501:1-12.
- Forbes-Osborne MA, Wilson SG, Morris AC. Insulinoma-associated 1a (Insm1a) is required for photoreceptor differentiation in the zebrafish retina. *Dev Biol.* 2013;380:157-171.
- Hori K, Sen A, Artavanis-Tsakonas S. Notch signaling at a glance. *J Cell Sci.* 2013;126:2135-2140.
- Artavanis-Tsakonas S, Rand MD, Lake RJ. Notch signaling: cell fate control and signal integration in development. *Science.* 1999;284:770-776.
- Bernardos RL, Lentz SI, Wolfe MS, Raymond PA. Notch-Delta signaling is required for spatial patterning and Müller glia differentiation in the zebrafish retina. *Dev Biol.* 2005;278:381-395.
- Jadhav AP, Cho S-H, Cepko CL. Notch activity permits retinal cells to progress through multiple progenitor states and acquire a stem cell property. *Proc Natl Acad Sci U S A.* 2006;103:18998-19003.
- Scheer N, Groth A, Hans S, Campos-Ortega JA. An instructive function for Notch in promoting gliogenesis in the zebrafish retina. *Development.* 2001;128:1099-1107.
- Uribe RA, Kwon T, Marcotte EM, Gross JM. Id2a functions to limit Notch pathway activity and thereby influence the transition from proliferation to differentiation of retinoblasts during zebrafish retinogenesis. *Dev Biol.* 2012;371:280-292.
- Perron M, Harris WA. Determination of vertebrate retinal progenitor cell fate by the Notch pathway and basic helix-loop-helix transcription factors. *Cell Mol Life Sci.* 2000;57:215-223.
- Nelson BR, Hartman BH, Ray CA, Hayashi T, Bermingham-McDonogh O, Reh TA. Acheate-scute like 1 (Ascl1) is required for normal delta-like (Dll) gene expression and notch signaling during retinal development. *Dev Dyn.* 2009;238:2163-2178.
- Nelson BR, Reh TA. Relationship between Delta-like and proneural bHLH genes during chick retinal development. *Dev Dyn.* 2008;237:1565-1580.
- Bernardos RL, Barthel LK, Meyers JR, Raymond PA. Late-stage neuronal progenitors in the retina are radial Müller glia that function as retinal stem cells. *J Neurosci.* 2007;27:7028-7040.
- Fausett BV, Goldman D. A role for alpha1 tubulin-expressing Müller glia in regeneration of the injured zebrafish retina. *J Neurosci.* 2006;26:6303-6313.
- Goldman D. Müller glial cell reprogramming and retina regeneration. *Nat Chem Biol.* 2014;1-12.
- Gorsuch RA, Hyde DR. Regulation of Müller glial dependent neuronal regeneration in the damaged adult zebrafish retina. *Exp Eye Res.* 2014;123:131-140.
- Lenkowski JR, Raymond PA. Müller glia: stem cells for generation and regeneration of retinal neurons in teleost fish. *Prog Retin Eye Res.* 2014;40:94-123.
- Nagashima M, Barthel LK, Raymond PA. A self-renewing division of zebrafish Müller glial cells generates neuronal progenitors that require N-cadherin to regenerate retinal neurons. *Development.* 2013;140:4510-4521.
- Fimbel SM, Montgomery JE, Burket CT, Hyde DR. Regeneration of inner retinal neurons after intravitreal injection of ouabain in zebrafish. *J Neurosci.* 2007;27:1712-1724.
- Karl MO, Reh TA. Regenerative medicine for retinal diseases: activating endogenous repair mechanisms. *Trends Mol Med.* 2010;16:193-202.
- Hitchcock P, Ochocinska M, Sieh A, Otterson D. Persistent and injury-induced neurogenesis in the vertebrate retina. *Prog Retin Eye Res.* 2004;23:183-194.
- Conner C, Ackerman KM, Lahne M, Hobgood JS, Hyde DR. Repressing notch signaling and expressing tnf are sufficient to mimic retinal regeneration by inducing Müller glial prolifera-

- tion to generate committed progenitor cells. *J Neurosci.* 2014; 34:14403–14419.
35. Nelson CM, Ackerman KM, O'Hayer P, Bailey TJ, Gorsuch RA, Hyde DR. Tumor necrosis factor- $\alpha$  is produced by dying retinal neurons and is required for Müller glia proliferation during zebrafish retinal regeneration. *J Neurosci.* 2013;33:6524–6539.
  36. Ramachandran R, Zhao X-F, Goldman D. Insm1a-mediated gene repression is essential for the formation and differentiation of Müller glia-derived progenitors in the injured retina. *Nat Cell Biol.* 2012;14:1013–1023.
  37. Wan J, Ramachandran R, Goldman D. HB-EGF is necessary and sufficient for Müller glia dedifferentiation and retina regeneration. *Dev Cell.* 2012;22:334–347.
  38. Gemberling M, Bailey TJ, Hyde DR, Poss KD. The zebrafish as a model for complex tissue regeneration. *Trends Genet.* 2013; 29:611–620.
  39. Thomas JL, Ochocinska MJ, Hitchcock PF, Thummel R. Using the Tg(nrd:egfp)/albino zebrafish line to characterize in vivo expression of neurod. *PLoS One.* 2012;7:e29128.
  40. Nasevicius A, Ekker SC. Effective targeted gene 'knockdown' in zebrafish. *Nat Genet.* 2000;26:216–220.
  41. Carmany-Rampey A, Moens CB. Modern mosaic analysis in the zebrafish. *Methods.* 2006;39:228–238.
  42. Alvarez-Delfin K, Morris AC, Snelson CD, et al. Tbx2b is required for ultraviolet photoreceptor cell specification during zebrafish retinal development. *Proc Natl Acad Sci U S A.* 2009; 106:2023–2028.
  43. Link BA, Fadool JM, Malicki J, Dowling JE. The zebrafish young mutation acts non-cell-autonomously to uncouple differentiation from specification for all retinal cells. *Development.* 2000; 127:2177–2188.
  44. Fadool JM. Development of a rod photoreceptor mosaic revealed in transgenic zebrafish. *Dev Biol.* 2003;258:277–290.
  45. Hwang WY, Fu Y, Reyon D, et al. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol.* 2013;31: 227–229.
  46. Westerfield M. Chapter 9: Molecular methods. In: *The Zebrafish Book A Guide for the Laboratory Use of Zebrafish (Danio rerio)*. Eugene, OR: University of Oregon Press; 2000.
  47. Geling A, Steiner H, Willem M, Bally-Cuif L, Haass C. A gamma-secretase inhibitor blocks Notch signaling in vivo and causes a severe neurogenic phenotype in zebrafish. *EMBO Rep.* 2002;3: 688–694.
  48. Kawakami K, Abe G, Asada T, et al. zTrap: zebrafish gene trap and enhancer trap database. *BMC Dev Biol.* 2010;10:105.
  49. Scheer N, Campos-Ortega JA. Use of the Gal4-UAS technique for targeted gene expression in the zebrafish. *Mech Devel.* 1999;80:153–158.
  50. Luo J, Uribe RA, Hayton S, Calinescu A-A, Gross JM, Hitchcock PF. Midkine-A functions upstream of Id2a to regulate cell cycle kinetics in the developing vertebrate retina. *Neural Dev.* 2012; 7:33.
  51. Barthel LK, Raymond PA. Subcellular localization of  $\alpha$ -tubulin and opsin mRNA in the goldfish retina using digoxigenin-labeled cRNA probes detected by alkaline phosphatase and HRP histochemistry. *J Neurosci Methods.* 1993;50:145–152.
  52. David R, Wedlich D. PCR-based RNA probes: a quick and sensitive method to improve whole mount embryo in situ hybridizations. *Biotech.* 2001;30:769–775.
  53. Obholzer N, Wolfson S, Trapani JG, et al. Vesicular glutamate transporter 3 is required for synaptic transmission in zebrafish hair cells. *J Neurosci.* 2008;28:2110–2118.
  54. Vihtelic TS, Hyde DR. Light-induced rod and cone cell death and regeneration in the adult albino zebrafish (*Danio rerio*) retina. *J Neurobiol.* 2000;44:289–307.
  55. Vihtelic TS, Soverly JE, Kassen SC, Hyde DR. Retinal regional differences in photoreceptor cell death and regeneration in light-lesioned albino zebrafish. *Exp Eye Res.* 2006;82:558–575.
  56. Taylor S, Chen J, Luo J, Hitchcock P. *Light-Induced Photoreceptor Degeneration in the Retina of the Zebrafish. Methods in Molecular Biology.* Totowa, NJ: Humana Press; 2012:247–254.
  57. Thomas JL, Nelson CM, Luo X, Hyde DR, Thummel R. Characterization of multiple light damage paradigms reveals regional differences in photoreceptor loss. *Exp Eye Res.* 2012; 97:105–116.
  58. Bernardos RL, Raymond PA. GFAP transgenic zebrafish. *Gene Expr Patterns.* 2006;6:1007–1013.
  59. Thummel R, Bailey TJ, Hyde DR. In vivo electroporation of morpholinos into the adult zebrafish retina. *J Vis Exp.* 2011; 27:e3603.
  60. Craig SEL, Thummel R, Ahmed H, Vasta GR, Hyde DR, Hitchcock PF. The zebrafish galectin Drgal1-I2 is expressed by proliferating Müller glia and photoreceptor progenitors and regulates the regeneration of rod photoreceptors. *Invest Ophthalmol Vis Sci.* 2010;51:3244–3252.
  61. Thummel R, Kassen SC, Montgomery JE, Enright JM, Hyde DR. Inhibition of Müller glial cell division blocks regeneration of the light-damaged zebrafish retina. *Dev Neurobiol.* 2008;68: 392–408.
  62. Chapouton P, Skupien P, Hesl B, et al. Notch activity levels control the balance between quiescence and recruitment of adult neural stem cells. *J Neurosci.* 2010;30:7961–7974.
  63. Stenkamp DL. Neurogenesis in the fish retina. *Int Rev Cytol.* 2007;259:173–224.
  64. Zou J, Lathrop KL, Sun M, Wei X. Intact retinal pigment epithelium maintained by nok is essential for retinal epithelial polarity and cellular patterning in zebrafish. *J Neurosci.* 2008; 28:13684–13695.
  65. Schmitt EA, Dowling JE. Early retinal development in the zebrafish, *Danio rerio*: light and electron microscopic analyses. *J Comp Neurol.* 1999;404:515–536.
  66. Chen J, Rattner A, Nathans J. The rod photoreceptor-specific nuclear receptor Nr2e3 represses transcription of multiple cone-specific genes. *J Neurosci.* 2005;25:118–129.
  67. Shen YC, Raymond PA. Zebrafish cone-rod (crx) homeobox gene promotes retinogenesis. *Dev Biol.* 2004;269:237–251.
  68. Kok FO, Shin M, Ni C-W, et al. Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev Cell.* 2015;32:97–108.
  69. Schulte-Merker S, Stainier DY. Out with the old, in with the new: reassessing morpholino knockdowns in light of genome editing technology. *Development.* 2014;141:3103–3104.
  70. Agathocleous M, Harris WA. From progenitors to differentiated cells in the vertebrate retina. *Annu Rev Cell Dev Biol.* 2009; 25:45–69.
  71. Dyer MA, Cepko CL. Regulating proliferation during retinal development. *Nat Rev Neurosci.* 2001;2:333–342.
  72. Reese BE. Development of the retina and optic pathway. *Vision Res.* 2011;51:613–632.
  73. Swaroop A, Kim D, Forrest D. Transcriptional regulation of photoreceptor development and homeostasis in the mammalian retina. *Nat Chem Biol.* 2010;11:563–576.
  74. Epstein DJ, McMahon AP, Joyner AL. Regionalization of Sonic hedgehog transcription along the anteroposterior axis of the mouse central nervous system is regulated by Hnf3-dependent and -independent mechanisms. *Development.* 1999;126:281–292.
  75. Nemer G, Nemer M. Transcriptional activation of BMP-4 and regulation of mammalian organogenesis by GATA-4 and -6. *Dev Biol.* 2003;254:131–148.

76. Ramachandran R, Zhao X-F, Goldman D. Ascl1a/Dkk/beta-catenin signaling pathway is necessary and glycogen synthase kinase-3beta inhibition is sufficient for zebrafish retina regeneration. *Proc Natl Acad Sci U S A*. 2011;108:15858-15863.
77. Xiao YQ, Freire-de-Lima CG, Schiemann WP, Bratton DL, Vandivier RW, Henson PM. Transcriptional and translational regulation of TGF-beta production in response to apoptotic cells. *J Immunol*. 2008;181:3575-3585.
78. Zhuang L, Hulin J-A, Gromova A, et al. Barx2 and Pax7 have antagonistic functions in regulation of wnt signaling and satellite cell differentiation. *Stem Cells*. 2014;32:1661-1673.
79. Beurdeley M, Spatazza J, Lee HHC, et al. Otx2 binding to perineuronal nets persistently regulates plasticity in the mature visual cortex. *J Neurosci*. 2012;32:9429-9437.
80. Lesaffre B, Joliot A, Prochiantz A, Volovitch M. Direct non-cell autonomous Pax6 activity regulates eye development in the zebrafish. *Neural Dev*. 2007;2:2.
81. Nishida A, Furukawa A, Koike C, et al. Otx2 homeobox gene controls retinal photoreceptor cell fate and pineal gland development. *Nat Neurosci*. 2003;6:1255-1263.
82. Noguchi H, Bonner-Weir S, Wei FY, Matsushita M, Matsumoto S. BETA2/NeuroD protein can be transduced into cells due to an arginine- and lysine-rich sequence. *Diabetes*. 2005;54:2859-2866.
83. Fortini ME. Gamma-secretase-mediated proteolysis in cell-surface-receptor signalling. *Nat Rev Mol Cell Biol*. 2002;3:673-684.
84. Jadhav AP, Mason HA, Cepko CL. Notch 1 inhibits photoreceptor production in the developing mammalian retina. *Development*. 2006;133:913-923.
85. Henrique D, Hirsinger E, Adam J, et al. Maintenance of neuroepithelial progenitor cells by Delta-Notch signalling in the embryonic chick retina. *Curr Biol*. 1997;7:661-670.
86. Joshi I, Minter LM, Telfer J, et al. Notch signaling mediates G1/S cell-cycle progression in T cells via cyclin D3 and its dependent kinases. *Blood*. 2009;113:1689-1698.
87. Cohen B, Shimizu M, Izrailit J, et al. Cyclin D1 is a direct target of JAG1-mediated Notch signaling in breast cancer. *Breast Cancer Res Treat*. 2009;123:113-124.
88. Das D, Lanner F, Main H, et al. Notch induces cyclin-D1-dependent proliferation during a specific temporal window of neural differentiation in ES cells. *Dev Biol*. 2010;348:153-166.
89. Ronchini C, Capobianco AJ. Induction of cyclin D1 transcription and CDK2 activity by Notchic: implication for cell cycle disruption in transformation by Notchic. *Mol Cell Biol*. 2001;21:5925-5934.
90. Yu Y, Zhang Y, Guan W, et al. Androgen receptor promotes the oncogenic function of overexpressed jagged1 in prostate cancer by enhancing cyclin B1 expression via Akt phosphorylation. *Mol Cancer Res*. 2014;12:830-842.
91. Liu Y, Shen Y, Rest JS, Raymond PA, Zack DJ. Isolation and characterization of a zebrafish homologue of the cone rod homeobox gene. *Invest Ophthalmol Vis Sci*. 2001;42:481-487.
92. Morris AC, Scholz TL, Brockerhoff SE, Fadool JM. Genetic dissection reveals two separate pathways for rod and cone regeneration in the teleost retina. *Dev Neurobiol*. 2008;68:605-619.
93. Brockerhoff SE, Fadool JM. Genetics of photoreceptor degeneration and regeneration in zebrafish. *Cell Mol Life Sci*. 2011;68:651-659.
94. Gao Z, Ure K, Ables JL, et al. NeuroD1 is essential for the survival and maturation of adult-born neurons. *Nat Neurosci*. 2009;12:1090-1092.
95. Gregg RG, Willer GB, Fadool JM, Dowling JE, Link BA. Positional cloning of the young mutation identifies an essential role for the Brahma chromatin remodeling complex in mediating retinal cell differentiation. *Proc Natl Acad Sci U S A*. 2003;100:6535-6540.
96. Laranjeiro R, Whitmore D. Transcription factors involved in retinogenesis are co-opted by the circadian clock following photoreceptor differentiation. *Development*. 2014;141:2644-2656.
97. Udolph G, Rath P, Chia W. A requirement for Notch in the genesis of a subset of glial cells in the Drosophila embryonic central nervous system which arise through asymmetric divisions. *Development*. 2001;128:1457-1466.