

Effect of Body Position on Episcleral Venous Pressure in Healthy Subjects

Nitika Arora,¹ Jay W. McLaren,¹ David O. Hodge,² and Arthur J. Sit¹

¹Department of Ophthalmology, Mayo Clinic, Rochester, Minnesota, United States

²Department of Health Sciences Research, Mayo Clinic, Jacksonville, Florida, United States

Correspondence: Arthur J. Sit, Department of Ophthalmology, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA; sit.arthur@mayo.edu.

Submitted: May 1, 2017

Accepted: September 5, 2017

Citation: Arora N, McLaren JW, Hodge DO, Sit AJ. Effect of body position on episcleral venous pressure in healthy subjects. *Invest Ophthalmol Vis Sci.* 2017;58:5151-5156. DOI:10.1167/iov.17-22154

PURPOSE. The mechanism of IOP change during a body position change is poorly understood. In this study, we investigated changes in episcleral venous pressure (EVP) between two body positions, sitting and inclined, and compared this with changes in IOP.

METHODS. This study was a prospective, comparative case series of 43 eyes of 24 healthy volunteers. IOP was measured using a pneumatometer in the seated position. EVP was then measured in a selected episcleral vein by using an automated, slit-lamp-mounted venomanometer. Thirty minutes later, the subject was placed in an inclined position with the neck extended and the head resting on the chin rest of the slit lamp. After 5 minutes, IOP and EVP in the same vein were remeasured. EVP in the inclined position was compared with EVP in the seated position, and the change in IOP was compared with the change in EVP. Statistical significance was determined using generalized estimating equation models.

RESULTS. Mean IOP increased from 11.4 ± 3.0 mm Hg (mean \pm SD) in the sitting position to 13.1 ± 3.4 mm Hg in the inclined position ($P < 0.001$). Mean EVP increased from 6.4 ± 1.4 mm Hg in the sitting position to 7.8 ± 1.7 mm Hg in the inclined position ($P < 0.001$). The postural rise in IOP was not different from the rise in EVP ($P = 0.18$).

CONCLUSIONS. In the inclined position, IOP and EVP are higher than they are in the sitting position. The posture-induced rise in IOP can be attributed to an increase in EVP.

Keywords: episcleral venous pressure, intraocular pressure, prone position

IOP is the most important risk factor for glaucoma and is the only variable that can currently be modified for the treatment of glaucoma. IOP is dynamic and can change rapidly on a timescale of seconds to minutes, depending on the stimulus. It is dependent on body position, increases as posture changes from sitting to supine, and increases more when a subject is inverted.¹⁻⁵

Postural changes could in theory affect several variables that change the steady-state IOP. At steady state, IOP is determined by aqueous humor production (Q), outflow facility (C), uveoscleral outflow (U), and episcleral venous pressure (EVP) as described by Goldmann and later modified

$$IOP = EVP + \frac{1}{C}(Q - U) \quad (1)$$

Steady-state IOP can be increased or decreased by changes in any of these variables.⁶ IOP can also be affected transiently by deformation of the globe during eye movements, blinks, forced displacement, and changing choroidal blood volume.

The contributions of these variables to posture-induced changes in steady-state IOP have been investigated. Carlson et al.⁷ investigated the relationship between IOP and rate of aqueous humor formation while subjects were alternated between an upright and inverted body position. They found that body inversion increased IOP by as much as 11 mm Hg but did not change the aqueous flow rate significantly. Outflow facility also did not change between sitting and supine positions in a study by Selvadurai et al.⁸ A few studies have looked at the variation of episcleral venous pressure with body position,⁹⁻¹⁴

and EVP increased in supine and inverted postures compared to standing posture. However, the methods used to determine EVP did not use objective measurement end points and may have been inconsistent with pressure in episcleral veins measured directly by cannulation.^{15,16} The effect of body position on EVP, and its relationship to postural IOP change, therefore remains uncertain.

Understanding the mechanisms of IOP changes with posture are important for optimizing the treatment of glaucoma throughout the 24-hour period. In particular, during the nocturnal period, IOP is significantly higher than the diurnal period and is related to the recumbent position during sleep.^{17,18} Therapies targeted toward the mechanisms of postural IOP change may be useful in stabilizing IOP fluctuations, although evidence to support the clinical importance of postural IOP fluctuations in glaucoma pathogenesis is currently limited.

In this study, we measured EVP in sitting and recumbent postures by using semiautomated objective methods and examined the relationship between the increase in IOP and the change in EVP induced by changing from the sitting to inclined positions.

MATERIALS AND METHODS

Twenty-four healthy volunteers, 15 male and 9 female, ranging in age from 24 to 73 years, were recruited from Mayo Clinic employees and students and local area residents. A medical history was obtained, and a complete dilated eye examination






FIGURE 1. Computerized episcleral venomanometer.

was performed. Participants were excluded if they had an IOP greater than 22 mm Hg, evidence of glaucomatous optic neuropathy or any other ocular pathology, or a history of ocular surgery or therapy. Subjects were also excluded if they had high myopia (-6 diopters [D] or more) or high hyperopia ($+2$ D or more). Subjects with chronic medical conditions (e.g., hypertension) were allowed to participate provided that their medical condition had been under good control over the preceding 12 months and they did not use β -adrenergic antagonists systemically. Subjects were also checked to be sure they could extend their neck while in an inclined position and maintain this posture for 7 to 10 minutes (the typical time required to obtain measurements in both eyes) as required by the protocol. On the day of the study, participants were asked to maintain a regular sleep schedule and avoid excess caffeine. Our study was conducted in accordance with the principles of the Declaration of Helsinki (1989) and was reviewed and approved by the Institutional Review Board of Mayo Clinic. All subjects gave written consent after discussion of the nature and possible risks of the study.

Measurement Techniques

IOP was measured in both eyes by using a pneumatometer (Model 30 Classic; Reichert, Inc., Depew, NY, USA) after instilling proparacaine 0.5% in each eye. EVP was measured by using a slit-lamp-mounted, computerized venomanometer as described previously (Fig. 1).¹⁶ Briefly, a clear flexible membrane was placed against the surface of an episcleral vein, which was identified as being straighter and deeper than conjunctival vessels. The pressure behind the membrane was automatically and linearly increased by a computer-controlled motor drive to compress the vein. During compression, a high-definition video camera captured images of the vein as it collapsed, and pressure in the chamber was monitored, synchronized, and recorded with each video frame. The pressure associated with the initiation of venous collapse was assumed to be equal to the venous pressure and was determined from the video recording by using a method described previously.¹⁶

Measurement Protocol

Subjects were seated in a standard ophthalmologic examination chair. In all measurements, IOP was measured in the right eye before the left eye, and then EVP was measured in the right eye before the left eye. EVP was measured in the superior or superotemporal quadrant, and the mean of four measurements on one vein in each eye was calculated and accepted as episcleral venous pressure.




FIGURE 2. Subjects were placed in an inclined position on a surgical bed with the neck extended and the head resting on the chin rest of the slit lamp. The angle of the head-torso axis was approximately 45° .

After measurements were completed with the subject in the seated position, the subject rested for 30 minutes and then assumed an inclined position on a surgery stretcher (Stryker Corp., Kalamazoo, MI, USA) with the neck extended and the chin resting on the chin rest of the slit lamp so that torso of the patient was at an angle of 45° to the ground (Fig. 2). After maintaining this posture for 5 minutes to allow for autonomic, postural, and hormonal changes to stabilize,^{19,20} IOP and EVP were remeasured in both the eyes.

Statistical Analysis

IOP and EVP in the seated position were compared with IOP and EVP in the inclined position. In addition, the change in IOP between positions (Δ IOP) was compared to the change in EVP (Δ EVP). Significance of all comparisons were determined by using generalized estimating equation models to account for possible correlation between eyes of the same subject. The relationship between Δ IOP and Δ EVP was examined by using Pearson correlation and significance was determined by using generalized estimating equation models. The differences between Δ IOP and Δ EVP were expressed as a function of their means according to the methods of Bland and Altman.²¹ The limits of agreement, which include approximately 95% of the differences, were the mean difference between Δ IOP and Δ EVP ± 2 SDs of the difference.²¹

RESULTS

Forty-three eyes of 24 volunteers were studied, including 15 females and 9 males, with ages ranging from 24 to 73 years (43.3 ± 15.8 years, mean \pm SD; Table). Five eyes were excluded from the study because of poor-quality images of the episcleral veins, rendering accurate measurement of EVP unfeasible.

Mean IOP increased from 11.4 ± 3.0 mm Hg in the sitting position to 13.1 ± 3.4 mm Hg in the inclined position ($P < 0.001$). Mean EVP increased from 6.4 ± 1.4 mm Hg in the sitting position to 7.8 ± 1.7 mm Hg in the inclined position ($P < 0.001$; Fig. 3A).

TABLE. Baseline Characteristics of Study Participants

	Age, y	CCT, μ m	IOP, mm Hg	Refractive Error, D
Mean \pm SD	40.4 ± 15.5	549 ± 34	12.0 ± 2.1	-1.2 ± 1.7
Range	19 to 73	481 to 599	8 to 15	-6.0 to 1.3




FIGURE 3. (A) Mean IOP in the inclined position was greater than mean IOP in the sitting position ($P < 0.001$). (B) Mean EVP in the inclined position was greater than mean EVP in the sitting position ($P < 0.001$).

The postural rise in EVP from sitting to inclined positions was 1.4 ± 1.7 mm Hg and was not significantly different from the postural rise in IOP of 1.7 ± 1.1 mm Hg ($P = 0.18$; Fig. 3B). The minimum detectable difference between the Δ IOP and the Δ EVP was 0.77 mm Hg ($\alpha = 0.05$, $\beta = 0.20$, $n = 43$ eyes).

The Δ IOP was not correlated with the Δ EVP ($r = 0.24$, $P = 0.1$; Fig. 4); the minimum detectable correlation coefficient between the two differences was $r = 0.40$ ($\alpha = 0.05$, $\beta = 0.20$, $n = 43$ eyes). The limits of agreement between Δ IOP and Δ EVP were from -3.2 to 3.8 mm Hg (Fig. 5).

DISCUSSION

IOP fluctuations have been implicated as potential risk factors for glaucoma progression.²² Postural fluctuations in IOP are

very common,²³ but the mechanisms of these IOP fluctuations have not been fully understood. From the modified Goldmann model of IOP (Equation 1), we would expect that EVP and IOP change in a 1:1 ratio if outflow facility, aqueous humor flow rate, and unconventional outflow rate remain constant. In our study, changing from the sitting to inclined position increased both IOP and EVP by a small amount and the lack of a difference between the Δ IOP and the Δ EVP suggests that IOP increased, at least in part, because EVP increased. Previous studies that investigated the concordance of postural changes in IOP and EVP have reported variable results. Friberg et al.¹¹ measured EVP by using a magnifying, portable transducer in supine and head down vertical position. For every 0.83 ± 0.21 mm Hg increase in EVP, IOP increased by 1 mm Hg. In contrast, Lavery and Kiel²⁴ reported that EVP and IOP responses to head




FIGURE 4. Limits of agreement between the change in EVP and the change in IOP between positions, according to Bland and Altman (reference). The mean difference was 0.30 ± 2.5 mm Hg, and the limits of agreement that contained 95% of the measurements were -3.2 and 3.8 mm Hg.




FIGURE 5. The change in IOP was not correlated with the change in EVP. The minimum detectable correlation was $r = 0.40$.

down tilt in rabbits were similar, with measurements obtained via direct cannulation. Similarly, Linnér et al.¹⁰ and Leith⁹ showed that both IOP and EVP increased by approximately 1 mm Hg on changing from sitting to supine positions, when they measured EVP by using variations of the pressure chamber technique described by Seidel in 1923.^{9,10} However, neither of these two studies reported the minimum detectable difference between the change in IOP and the change in EVP, and both used subjective rather than objective measurement end points. In our study, we had an 80% chance of finding a difference as small as 0.77 mm Hg if it existed. If a smaller difference between the postural change in IOP and EVP existed, our study would not have been able to detect it. However, to detect a smaller difference between changes in these variables would be impractical; detecting a mean difference as small as 0.5 mm Hg would require 124 eyes.

Although other studies have evaluated the change in EVP with body position, we are not aware of any previous study that has evaluated EVP changes using an objective technique in living human eyes. Noninvasive measurement of EVP is based on identifying the pressure needed to collapse an episcleral vein to a specified end point.²⁵ However, only one end point is reflective of the correct EVP. Ideal Tube laws indicate that if a collapsible tube is subjected to an external pressure, the tube begins to collapse when the internal (venous) pressure is equal to the external (bulb) pressure.²⁶ Beyond this initial point of collapse, the pressure in the bulb can be much higher than the initial venous pressure. This was validated for EVP measurements in a study by Gaasterland et al.,¹⁵ in which noninvasive measurements obtained by a manual pressure chamber technique (analogous to the technique used in our device) was compared with measurements obtained by direct cannulation of the vessel. Selection of end points beyond the beginning point of collapse resulted in pressure measurements that could be more than two times greater than the true EVP. However, identification of the point of initial compression is difficult when performing measurements in conscious human subjects. The resulting lack of precision associated with using a subjective end point can make EVP measurements unreliable. Nevertheless, our study appears to confirm the results from previous studies using subjective techniques for EVP measurement in humans.

Potential differences between the postural IOP and EVP changes could result from changes in other parameters of the modified Goldmann equation. Although this study did not investigate the effect of body position on other parameters, previous studies^{7,8} reported that aqueous humor flow and outflow facility do not change between the recumbent and sitting positions. It is unknown if uveoscleral (unconventional) flow varies with body position. Also, the modified Goldmann equation is only valid for steady-state conditions, and any transient changes in IOP or other variables during the change in posture would invalidate our comparison between Δ IOP and Δ EVP. However, we allowed a minimum of 5 minutes after the change in body position to allow IOP and EVP to reach steady state in the new position. Other studies of IOP change with body position have reported that IOP increases immediately on assuming a headstand position and reaches a new steady state within 1 to 2 minutes.²⁷ The changes in body position in our study were much more limited, and we assumed that IOP would equilibrate just as rapidly. Whether EVP behaves in a similar manner has not been investigated, but adjustments in systemic cardiovascular parameters due to body position changes appear to occur within 30 seconds.²⁸

If postural IOP change was caused by postural EVP change alone, one would expect a correlation between these variables based on the modified Goldmann equation. The lack of a

significant correlation among our measurements was likely because of variability and the relatively small changes in these pressures between the sitting and supine positions. The correlation may have been better demonstrated if EVP had been elevated by a greater amount, for example, by inversion. Unfortunately, our method for measuring EVP was not suited for measurements in an inverted subject. Another possibility is that the measurements of IOP and EVP simply represent two time points. Although we performed the measurements of EVP and IOP as close together as possible, they are not simultaneous measurements, and these parameters can vary relatively quickly with changes in intra-abdominal pressure or even deep respiration. This limitation highlights the importance of interpreting population differences instead of individual results in aqueous humor dynamics studies using current technology. Finally, it is possible that changes in IOP and EVP are temporally associated but vary due to independent causes yet to be identified.

In glaucoma patients, it is unknown if postural EVP changes, as well as the relationship between IOP and EVP, are similar to what we found in this study of normal subjects. Reported postural changes in IOP from sitting to supine in normal subjects range from 0.3 to 5.6 mm Hg or more, depending on the study, whereas, in glaucoma patients, the reported change is greater, ranging from 1.6 mm Hg to more than 8.6 mm Hg.^{5,18} Whether EVP is altered in glaucoma (other than in Sturge-Weber or carotid cavernous fistulas) is unclear, and various studies have reported conflicting results.²⁹⁻³³ If postural changes in IOP are directly related to postural changes in EVP in glaucoma patients, then we would expect to see a greater elevation in inclined EVP compared with normal eyes. However, further measurements in glaucomatous eyes are needed to understand changes in IOP and EVP with postural variations.

The mechanism for EVP change with posture is likely related to changes in hydrostatic pressure. However, the magnitude of EVP change is far less than would be expected from a simple fluid column. Assuming a distance of 20 cm between the eyes and the heart, we would expect a hydrostatic pressure difference of 14.7 mm Hg between the upright and recumbent positions if the venous system was an open channel. In the 45° position of our study, this would still result in a change in height of 14.1 cm or 10.4 mm Hg in hydrostatic pressure. Clearly this does not occur because orthostatic changes in venous pressures are regulated by the sympathetic system.³⁴ Larger transient changes in pressure may occur, but measurements in our study were performed after at least 5 minutes after changing positions, whereas the orthostatic response in blood pressure is complete in less than 1 minute.³⁵ Another regulatory mechanism may stem from the possibility that intraocular veins form a Starling resistor because they are subject to the surrounding IOP, which would prevent the venous pressure from dropping below IOP.³⁶ Although episcleral veins measured in this study were relatively superficial, they are part of a venous plexus that includes intrascleral and intraocular vessels.

A limitation of our study is the technique used to measure EVP in the inclined position. In the 45° position used in our study, subjects were semiprone, and their head and neck were elevated to accommodate the slit lamp (Fig. 2). This positioning was necessary with the current slit-lamp-based equipment used to measure EVP, which limits the head position to upright. The slit lamp is critical for EVP measurements several reasons. First, it provides a method to visualize the episcleral veins of interest. These vessels are typically 50 to 100 μ m in diameter and, without sufficient magnification and illumination, it would be difficult to differentiate between episcleral veins and other superficial

vessels. Adequate magnification is also required to obtain images of sufficient resolution to detect changes in brightness easily. As well, the slit lamp provides an important point of stabilization at high magnification, enabling the capture of high quality videos. Although our long-term goals include the design and construction of a completely new system enabling EVP measurements in any position, this is far beyond the scope of the current study.

The changes in IOP and EVP in our study may have been greater if subjects had been positioned completely horizontal or had been inverted. Alternately, the change in EVP and IOP could have been elevated by the extended head and neck position, which has been reported to affect IOP.²³ It is not known if the relationship between EVP and IOP is affected by alterations of the head and neck position. However, if conditions are at steady state at the time of measurements, the Goldmann equation should be valid. Nevertheless, further investigation of the elevation of EVP and its contribution to the elevation of IOP in the supine position will need to wait for the development of smaller and more flexible instruments that permit measurements of EVP in any position.

Another potential limitation is that EVP measurements from our device have not been compared with venous pressures measured by direct cannulation. However, as discussed above, a study by Gaasterland et al.¹⁵ compared noninvasive EVP measurements using a manual pressure chamber technique analogous to our device with invasive measurements in anesthetized monkeys and found that the earliest detectable vein collapse most closely represented EVP measured by cannulation. However, this end point is extremely difficult to identify in real time when performing measurements in conscious human subjects. Our technique identified the initial point of collapse using image analysis of videos and associated pressure measurements during automated episcleral vein compression.

In summary, both IOP and EVP are higher in the inclined position than they are in the sitting position. Our study suggests that postural changes in IOP are consistent with the postural changes in EVP, as indicated by the Goldmann equation. Further investigation is required to determine the contribution of EVP to nonpostural IOP fluctuations. As well, further investigation is required to understand the role of modulating EVP to reduce IOP variability.

Acknowledgments

Supported by the Bright Focus Foundation, Mayo Foundation for Medical Education and Research, and an unrestricted departmental grant from Research to Prevent Blindness.

Disclosure: **N. Arora**, None; **J.W. McLaren**, None; **D.O. Hodge**, None; **A.J. Sit**, None

References

- Ozcan MS, Praetel C, Bhatti MT, Gravenstein N, Mahla ME, Seubert CN. The effect of body inclination during prone positioning on intraocular pressure in awake volunteers: a comparison of two operating tables. *Anesth Analg*. 2004;99:1152-1158.
- Walick KS, Kragh JE Jr, Ward JA, Crawford JJ. Changes in intraocular pressure due to surgical positioning: studying potential risk for postoperative vision loss. *Spine (Phila Pa 1976)*. 2007;32:2591-2595.
- Cheng MA, Todorov A, Tempelhoff R, McHugh T, Crowder CM, Laurysen C. The effect of prone positioning on intraocular pressure in anesthetized patients. *Anesthesiology*. 2001;95:1351-1355.
- Grant GP, Szirth BC, Bennett HL, et al. Effects of prone and reverse trendelenburg positioning on ocular parameters. *Anesthesiology*. 2010;112:57-65.
- Prata TS, De Moraes CG, Kanadani FN, Ritch R, Paranhos A Jr. Posture-induced intraocular pressure changes: considerations regarding body position in glaucoma patients. *Surv Ophthalmol*. 2010;55:445-453.
- Goldmann H. Out-flow pressure, minute volume and resistance of the anterior chamber flow in man. *Doc Ophthalmol*. 1951;5-6:278-356.
- Carlson KH, McLaren JW, Topper JE, Brubaker RE. Effect of body position on intraocular pressure and aqueous flow. *Invest Ophthalmol Vis Sci*. 1987;28:1346-1352.
- Selvadurai D, Hodge D, Sit AJ. Aqueous humor outflow facility by tonography does not change with body position. *Invest Ophthalmol Vis Sci*. 2010;51:1453-1457.
- Leith AB. Episcleral venous pressure in tonography. *Br J Ophthalmol*. 1963;47:271-278.
- Linnér E, Rickenbach C, Werner H. Comparative measurements of the pressure in the aqueous veins and the conjunctival veins using different methods. *Acta Ophthalmol (Copenb)*. 1950;28:469-478.
- Friberg TR, Sanborn G, Weinreb RN. Intraocular and episcleral venous pressure increase during inverted posture. *Am J Ophthalmol*. 1987;103:523-526.
- Friberg TR, Weinreb RN. Ocular manifestations of gravity inversion. *JAMA*. 1985;253:1755-1757.
- Blondeau P, Tetrault JP, Papamarkakis C. Diurnal variation of episcleral venous pressure in healthy patients: a pilot study. *J Glaucoma*. 2001;10:18-24.
- Sultan M, Blondeau P. Episcleral venous pressure in younger and older subjects in the sitting and supine positions. *J Glaucoma*. 2003;12:370-373.
- Gaasterland DE, Pederson JE. Episcleral venous pressure: a comparison of invasive and noninvasive measurements. *Invest Ophthalmol Vis Sci*. 1983;24:1417-1422.
- Sit AJ, Ekdawi NS, Malihi M, McLaren JW. A novel method for computerized measurement of episcleral venous pressure in humans. *Exp Eye Res*. 2011;92:537-544.
- Liu JH, Zhang X, Kripke DE, Weinreb RN. Twenty-four-hour intraocular pressure pattern associated with early glaucomatous changes. *Invest Ophthalmol Vis Sci*. 2003;44:1586-1590.
- Nau CB, Malihi M, McLaren JW, Hodge DO, Sit AJ. Circadian variation of aqueous humor dynamics in older healthy adults. *Invest Ophthalmol Vis Sci*. 2013;54:7623-7629.
- Jain MR, Marmion VJ. Rapid pneumatic and Mackey-Marg applanation tonometry to evaluate the postural effect on intraocular pressure. *Br J Ophthalmol*. 1976;60:687-693.
- Rössler A, Laszlo Z, Haditsch B, Hinghofer-Szalkay HG. Orthostatic stimuli rapidly change plasma adrenomedullin in humans. *Hypertension*. 1999;34:1147-1151.
- Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. [see comment]. *Lancet*. 1986;1:307-310.
- Sit AJ. Intraocular pressure variations: causes and clinical significance. *Can J Ophthalmol*. 2014;49:484-488.
- Malihi M, Sit AJ. Effect of head and body position on intraocular pressure. *Ophthalmology*. 2012;119:987-991.
- Lavery WJ, Kiel JW. Effects of head down tilt on episcleral venous pressure in a rabbit model. *Exp Eye Res*. 2013;111:88-94.
- Sit AJ, McLaren JW. Measurement of episcleral venous pressure. *Exp Eye Res*. 2011;93:291-298.
- Shapiro AH. Steady flow in collapsible tubes. *J Biomech Eng*. 1977;99:126-147.

27. Jasien JV, Jonas JB, de Moraes CG, Ritch R. Intraocular pressure rise in subjects with and without glaucoma during four common yoga positions. *PLoS One*. 2015;10:e0144505.
28. Toska K, Walloe L. Dynamic time course of hemodynamic responses after passive head-up tilt and tilt back to supine position. *J Appl Physiol (1985)*. 2002;92:1671-1676.
29. Podos SM, Minas TF, Macri FJ. A new instrument to measure episcleral venous pressure. Comparison of normal eyes and eyes with primary open-angle glaucoma. *Arch Ophthalmol*. 1968;80:209-213.
30. Talusan ED, Schwartz B. Episcleral venous pressure. Differences between normal, ocular hypertensive, and primary open angle glaucomas. *Arch Ophthalmol*. 1981;99:824-828.
31. Weigelin E, Lohlein H. Measurements of the blood pressure of the episcleral vessels of the eye in healthy persons. *Albrecht Von Graefes Arch Ophthalmol*. 1952;153:202-213.
32. Selbach JM, Posielek K, Steuhl KP, Kremmer S. Episcleral venous pressure in untreated primary open-angle and normal-tension glaucoma. *Ophthalmologica*. 2005;219:357-361.
33. Johnson TV, Fan S, Camras CB, Toris CB. Aqueous humor dynamics in exfoliation syndrome. *Arch Ophthalmol*. 2008;126:914-920.
34. Stewart JM. Mechanisms of sympathetic regulation in orthostatic intolerance. *J Appl Physiol (1985)*. 2012;113:1659-1668.
35. Wieling W, Krediet CT, van Dijk N, Linzer M, Tschakovsky ME. Initial orthostatic hypotension: review of a forgotten condition. *Clin Sci (Lond)*. 2007;112:157-165.
36. Kiel JW. The ocular circulation. In: Granger DN, Granger JP, eds. *Integrated Systems Physiology: From Molecule to Function Disease*. San Rafael, CA: Morgan & Claypool Life Sciences; 2010.